

Phase transition on Exel crossed products associated to dilation matrices

MARCELO LACA IAIN RAEBURN

JACQUI RAMAGGE*

Marcelo Laca, Department of Mathematics and Statistics
University of Victoria
Victoria, BC V8W 3P4
Canada

Iain Raeburn, Department of Mathematics and Statistics
University of Otago
PO Box 56
Dunedin 9054
New Zealand

Jacqui Ramagge, School of Mathematics and Applied Statistics
University of Wollongong
NSW 2522
Australia

Email: laca@math.uvic.ca

Email: iraeburn@maths.otago.ac.nz

Email: ramagge@uow.edu.au

25 January 2011

ABSTRACT. An integer matrix $A \in M_d(\mathbb{Z})$ induces a covering σ_A of \mathbb{T}^d and an endomorphism $\alpha_A: f \mapsto f \circ \sigma_A$ of $C(\mathbb{T}^d)$ for which there is a natural transfer operator L . In this paper, we compute the KMS states on the Exel crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ and its Toeplitz extension. We find that $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ has a unique KMS state, which has inverse temperature $\beta = \log |\det A|$. Its Toeplitz extension, on the other hand, exhibits a phase transition at $\beta = \log |\det A|$, and for larger β the simplex of KMS_β states is isomorphic to the simplex of probability measures on \mathbb{T}^d .

1. INTRODUCTION

Actions of the real line \mathbb{R} on C^* -algebras are used to describe the time evolution in physical models, and also arise in a wide variety of mathematical contexts. The KMS states for the action were originally intended to be mathematical realisations of the equilibrium states in statistical mechanics

*. This research was supported by the Natural Sciences and Engineering Research Council of Canada, by the University of Otago, and by the Australian Research Council.

[bra-rob]. More recently, mathematicians have found actions of \mathbb{R} on algebras of number-theoretic origin that exhibit phase transitions of the sort one might expect in a statistical-mechanical model [bos-con, laca98, lr]. Here we describe a similar phenomenon for the gauge action on an Exel crossed-product C^* -algebra associated to an integer dilation matrix A .

An illuminating example for the analysis of KMS states is the action σ lifted from the gauge action of \mathbb{T} on the Toeplitz-Cuntz algebra \mathcal{TO}_n [eva]. The system (\mathcal{TO}_n, σ) has a single KMS state for each inverse temperature $\beta \geq \log n$, but only the one at $\beta = \log n$ factors through the purely infinite simple quotient \mathcal{O}_n (see, for example, [ln, Example 2.8]). Our situation is similar: the Exel crossed product is purely infinite simple and has a unique KMS state, which has inverse temperature $\beta = \log |\det A|$, whereas its Toeplitz analogue has KMS_β states for all $\beta \geq \log |\det A|$. Here, though, the simplex of KMS_β states is large for $\beta > \log |\det A|$, and we have a phase transition at $\beta = \log |\det A|$.

Before stating our results more precisely, we set up some notation. We consider a matrix $A \in M_d(\mathbb{Z})$ with nonzero determinant, and write σ_A for the associated self-covering of $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$. Then $\alpha_A: f \mapsto f \circ \sigma_A$ is an endomorphism of $C(\mathbb{T}^d)$,

$$L(f)(z) = \frac{1}{|\det A|} \sum_{\sigma_A(w)=z} f(w) \quad (1)$$

defines a transfer operator L for α_A , and the triple $(C(\mathbb{T}^d), \alpha_A, L)$ is one of the Exel systems studied in [ehr]. We write M_L for the associated right-Hilbert bimodule over $C(\mathbb{T}^d)$, which has underlying space $C(\mathbb{T}^d)$, actions defined by $f \cdot m \cdot g = f m \alpha_A(g)$, and inner product defined by $\langle m, n \rangle = L(m^* n)$. We write ϕ for the homomorphism of $C(\mathbb{T}^d)$ into $\mathcal{L}(M_L)$ which implements the left action.

If Σ is a set of coset representatives for $\mathbb{Z}^d / A^t \mathbb{Z}^d$, then the characters $\{\gamma_m: z \mapsto z^m : m \in \Sigma\}$, viewed as continuous functions on \mathbb{T}^d and hence as elements of M_L , form an orthonormal basis for M_L (this observation is due to Packer and Rieffel [pr], and a proof consistent with our notation is given in [ehr, Lemma 2.6]). The reconstruction formula for this basis implies that $\phi(f)$ is the finite-rank operator $\sum_m \Theta_{f \cdot \gamma_m, \gamma_m}$ for every $f \in C(\mathbb{T}^d)$. Then since α_A is unital, the results of [br] imply that Exel's Toeplitz algebra $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L)$ is the Toeplitz algebra $\mathcal{T}(M_L)$, and that the Exel crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ is isomorphic to the Cuntz-Pimsner algebra $\mathcal{O}(M_L)$.

The Toeplitz algebra $\mathcal{T}(M)$ of a Hilbert bimodule M over C is generated by a universal representation (i_M, i_C) of M , and carries a gauge action of \mathbb{T} characterised by $\gamma_z(i_M(m)) = z i_M(m)$ and $\gamma_z(i_C(c)) = i_C(c)$; this action descends to the Cuntz-Pimsner algebra $(\mathcal{O}(M), j_M, j_C)$. The gauge actions inflate to actions σ of \mathbb{R} which are characterised by

$$\begin{aligned} \sigma_t \circ i_C &= i_C \text{ and } \sigma_t(i_M(m)) = e^{it} i_M(m), \text{ and} \\ \sigma_t \circ j_C &= j_C \text{ and } \sigma_t(j_M(m)) = e^{it} j_M(m). \end{aligned}$$

Our goal is the following description of the KMS states of $(\mathcal{T}(M_L), \sigma)$ and $(\mathcal{O}(M_L), \sigma)$.

Theorem 1. *Suppose that $A \in M_d(\mathbb{Z})$ has nonzero determinant, $(C(\mathbb{T}^d), \alpha_A, L)$ is the associated Exel system, and σ denotes the action of \mathbb{R} on $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L)$ satisfying (1).*

(a) *There are no KMS_β states on $(\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L), \sigma)$ unless $\beta \geq \log |\det A|$.*

(b) *For each $\beta \in (\log |\det A|, \infty]$, the simplex of KMS_β states on $(\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L), \sigma)$ is affinely homeomorphic to the simplex $P(\mathbb{T}^d)$ of probability measures on \mathbb{T}^d .*

(c) *If A is a dilation matrix, then $(\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L), \sigma)$ has a unique $\text{KMS}_{\log |\det A|}$ state, and this state factors through the quotient map*

$$Q: (\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L), \sigma) \rightarrow (C(\mathbb{T}^d) \rtimes_{\alpha, L} \mathbb{N}, \sigma).$$

(d) *Every ground state of $(\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L), \sigma)$ is a KMS_∞ state.*

After a short review of notation and conventions, we begin in §3 by giving presentations of $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L)$ and $C(\mathbb{T}^d) \rtimes_{\alpha, L} \mathbb{N}$ in terms of a unitary representation u of \mathbb{Z}^d and an isometry v which, loosely, implements the action α_A . Then in §4, we characterise the KMS states of $(\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L), \sigma)$ in terms of their behaviour with respect to the presentation in §3.

We then set about proving Theorem 1 in stages, and we give more precise formulations of our results as we go. For example, we prove part (c) in §5, and we prove a little more than we stated above: we only need to assume that A is a dilation matrix to get uniqueness of the $\text{KMS}_{\log |\det A|}$ state. In §6, we prove existence of lots of KMS states (see Proposition 18); a novelty in our construction is the use of induced representations to build Hilbert spaces where we can construct KMS states from vector states. Then in §7, we prove that we have found all the KMS_β states for $\beta > \log |\det A|$. In §8, we prove part (d).

Theorem 1 and our strategy for proving it were motivated by our previous work in [lr], or more precisely, by what it says about the KMS states of the additive boundary quotient $(\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes \mathbb{N}^\times), \sigma)$ of the Toeplitz algebra $\mathcal{T}(\mathbb{N} \rtimes \mathbb{N}^\times)$ (see [BaHLR, §4]). The connection with Exel crossed products is made in [BaHLR, §5], where it is shown that there is an Exel system $(C(\mathbb{T}), \alpha, L, \mathbb{N}^\times)$ of the kind studied in [l] whose Nica-Toeplitz crossed product $\mathcal{NT}(C(\mathbb{T}), \alpha, L, \mathbb{N}^\times)$ is $\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes \mathbb{N}^\times)$ and whose Exel crossed product $C(\mathbb{T}) \rtimes_{\alpha, L} \mathbb{N}^\times$ is the Crisp-Laca boundary quotient of $\mathcal{T}(\mathbb{N} \rtimes \mathbb{N}^\times)$ (or in other words, Cuntz's $\mathcal{Q}_\mathbb{N}$ [c2]). So our present analysis differs from that in [lr] in that we have raised the dimension of the torus to d , but have replaced $\mathbb{N}^\times \cong \mathbb{N}^\infty$ by \mathbb{N} . The case $d=1$, where A has the form (N) , is in some sense an intersection of our results with those in [lr], and in §9 we carry out an analysis of the KMS states on $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ parallel to that in [BaHLR, §4].

We close in §10 with a discussion of the case where A is invertible over \mathbb{Z} . The endomorphism α_A is then an automorphism, and $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ is the usual crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$, which can also be viewed as a group algebra $C^*(\mathbb{Z}^d \rtimes_{A^t} \mathbb{Z})$. We know from §5 that there can only be KMS_β states when $\beta = \log |\det A| = 0$, so we are left to determine the invariant traces, which we do in Proposition 32. When $\mathbb{Z}^d \rtimes_{A^t} \mathbb{Z}$ is the integer Heisenberg group, for example, we can find lots of invariant traces.

2. NOTATION AND COVENTIONS

2.1. Integer matrices..

Throughout this paper A is a matrix in $M_d(\mathbb{Z})$ whose determinant $\det A$ is nonzero. If the eigenvalues λ of $A \in M_d(\mathbb{Z})$ all satisfy $|\lambda| > 1$, then we call A a *dilation matrix*. This was a standing assumption in [ehr], but here we do not in general assume that A is a dilation matrix. We use multiindex notation, so that $e^{2\pi i x} = (e^{2\pi i x_1}, \dots, e^{2\pi i x_d})$ for $x \in \mathbb{R}^d$, and the covering map $\sigma_A: \mathbb{T}^d \rightarrow \mathbb{T}^d$ is characterised by $\sigma_A(e^{2\pi i x}) = e^{2\pi i Ax}$ for $x \in \mathbb{R}^d$. Since the transpose A^t appears more often than A , we write $B := A^t$; we have tried to avoid using the letters A and B for anything else. We choose a set Σ of coset representatives for $\mathbb{Z}^d / B\mathbb{Z}^d$, and assume for convenience that $0 \in \Sigma$. We sometimes write N for $|\det A| = |\det B|$.

2.2. Hilbert bimodules.

A bimodule M over a C^* -algebra C is a right-Hilbert bimodule if it is a right Hilbert C -module, and if the left action of C is implemented by a homomorphism ϕ of C into the C^* -algebra $\mathcal{L}(M)$ of adjointable operators. (Such bimodules are also called “correspondences” over C , or just “Hilbert bimodules” for short.) Our C^* -algebras will always have identities, and our bimodules are always essential in the sense that $\phi: C \rightarrow \mathcal{L}(M)$ is unital.

A representation¹ (ψ, π) of a Hilbert bimodule M in a C^* -algebra D consists of a linear map $\psi: M \rightarrow D$ and a unital representation $\pi: C \rightarrow D$ such that

$$\psi(c_1 \cdot m \cdot c_2) = \pi(c_1) \psi(m) \pi(c_2) \text{ and } \pi(\langle m, n \rangle) = \psi(m)^* \psi(n).$$

Every Hilbert bimodule M has a Toeplitz algebra $\mathcal{T}(M)$, which is generated by a universal representation (i_M, i_C) .

Every representation (ψ, π) of M in D induces a representation $(\psi, \pi)^{(1)}: \mathcal{K}(M) \rightarrow D$ such that $(\psi, \pi)^{(1)}(\Theta_{m,n}) = \psi(m) \psi(n)^*$ (see [p, page 202] or [fr, Proposition 1.6]). The representation (ψ, π) is Cuntz-Pimsner covariant if

$$(\psi, \pi)^{(1)}(\phi(a)) = \pi(a) \text{ whenever } \phi(a) \in \mathcal{K}(M).$$

1. These are often called Toeplitz representations, but we now believe this to have been an unfortunate choice of name (see [BaHLR, Remark 5.3]).

The Cuntz-Pimsner algebra $\mathcal{O}(M)$ is the quotient of $\mathcal{T}(M)$ that is universal for Cuntz-Pimsner covariant representations. We write $Q: \mathcal{T}(M) \rightarrow \mathcal{O}(M)$ for the quotient map, and $(j_M, j_C) := (Q \circ i_M, Q \circ i_C)$ for the universal Cuntz-Pimsner covariant representation in $\mathcal{O}(M)$. (Though there are several different definitions of Cuntz-Pimsner covariance out there, they all coincide for the bimodules in this paper.)

2.3. Exel crossed products.

An Exel system consists of an endomorphism α of a C^* -algebra C , and a transfer operator L for α , which is a bounded positive linear map $L: C \rightarrow C$ such that $L(\alpha(c)d) = cL(d)$. The examples of interest here are the systems $(C(\mathbb{T}^d), \alpha_A, L)$ discussed in the introduction, where α_A is the endomorphism $f \mapsto f \circ \sigma_A$ associated to an integer matrix A , and L is defined by averaging over inverse images of points, as in (1). Notice that both α and L are unital.

Every Exel system (C, α, L) gives rise to a Hilbert bimodule over C as follows. We first make a copy C_L of C into a bimodule over C by setting $c \cdot m = cm$ and $m \cdot c = m\alpha(c)$ for $m \in C_L$ and $c \in C$. The formula $\langle m, n \rangle := L(m^*n)$ carries a C -valued pre-inner product on C_L , and completing C_L gives a right Hilbert C -module M_L . Because L is bounded, the left action of C extends to an action of C by adjointable operators on the completion M_L . (The details are in [br, §3].) In general the completion process involves modding out by vectors of length zero, so that the quotient carries a C -valued inner product. However, for the systems $(C(\mathbb{T}^d), \alpha_A, L)$, the module $C(\mathbb{T}^d)$ has no vectors of length zero and is already complete (see [LR2, Lemma 3.3]). So we dispense with the quotient maps $q: C_L \rightarrow M_L$ which were used in [br] to distinguish between elements of the algebra and elements of the bimodule.

For an Exel system (C, α, L) , we define the Toeplitz algebra $\mathcal{T}(C, \alpha, L)$ to be $\mathcal{T}(M_L)$, and the Exel crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ to be the Cuntz-Pimsner algebra $\mathcal{O}(M_L)$. This is not quite Exel's original definition [e1], but for the systems $(C(\mathbb{T}^d), \alpha_A, L)$ of interest to us it is equivalent. (The precise relationship between Exel's crossed product and Cuntz-Pimsner algebras is worked out in [br, §3].) So $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L)$ and $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ are generated by universal representations $(i_{M_L}, i_{C(\mathbb{T}^d)})$ and $(j_{M_L}, j_{C(\mathbb{T}^d)})$.

2.4. KMS states.

Suppose that σ is an action of \mathbb{R} by automorphisms of a C^* -algebra C . An element c of C is analytic if $t \mapsto \sigma_t(c)$ is the restriction of an entire function. A state ϕ of C is a KMS state at inverse temperature $\beta \in (0, \infty)$ if there is a set S of analytic elements such that $\text{span} S$ is dense in C and

$$\phi(dc) = \phi(c\sigma_{it}(d)) \text{ for } c, d \in S.$$

In [lr, §7], we were careful to explain why this definition is equivalent to that used in the standard sources [bra-rob] and [ped]. We also adopt two more recent conventions which are possibly nonstandard. First, we regard the KMS_0 states to be the σ -invariant traces; this agrees with the convention in [ped] rather than the one in [bra-rob]. Second, we use the conventions

of Connes and Marcolli [cm], which distinguish between the KMS_∞ states (those which are weak* limits of KMS_β states as $\beta \rightarrow \infty$) and the ground states (those such that $z \mapsto \phi(c\sigma_z(d))$ is bounded on the upper half-plane). Neither [bra-rob] nor [ped] makes this distinction.

3. A PRESENTATION

We describe a presentation of the Toeplitz algebra $\mathcal{T}(M_L)$ like that of $\mathcal{T}(\mathbb{N} \rtimes \mathbb{N}^\times)$ in [lr, Theorem 4.1], or, more precisely, like that of the additive boundary quotient $\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes \mathbb{N}^\times)$ in [BaHLR, Proposition 3.3].

Proposition 2. *Suppose that $A \in M_d(\mathbb{Z})$ has $\det A \neq 0$, and consider the Exel system $(C(\mathbb{T}^d), \alpha_A, L)$. Then the Toeplitz algebra $\mathcal{T}(M_L)$ is the universal C^* -algebra generated by a unitary representation $u: \mathbb{Z}^d \rightarrow U(\mathcal{T}(M_L))$ and an isometry $v \in \mathcal{T}(M_L)$ satisfying*

$$(E1) \quad v u_m = u_{Bm} v, \text{ and}$$

$$(E2) \quad v^* u_m v = \begin{cases} u_{B^{-1}m} & \text{if } m \in B\mathbb{Z}^d \\ 0 & \text{otherwise.} \end{cases}$$

If U is a unitary representation of \mathbb{Z}^d in a C^* -algebra C and $V \in C$ is an isometry satisfying (E1) and (E2), then the corresponding representation (ψ, π) of M_L satisfies $U_m = \pi(\gamma_m)$ and $V = \psi(1)$.

Remark 3. *It might be helpful to see how our presentation is related to the presentation of $\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes \mathbb{N}^\times)$ in [BaHLR]. The isometries $\{v_p: p \in \mathcal{P}\}$ have become the single isometry v , and the additive generator s has been replaced by the unitary representation u of the additive group \mathbb{Z}^d . The relations (T2) and (T3) in [BaHLR, Proposition 3.3] are not needed because here “we only have one prime” (which we will normalise to e when we define our dynamics!), and the relation (Q6) in [BaHLR, Proposition 3.3] is replaced by the assumption that u is a unitary representation. So we are left with (T1) and (T5), which are analogous to (E1) and (E2) respectively.*

The relation (E2) implies that $\{u_m v: m \in \Sigma\}$ is a Toeplitz-Cuntz family. The analogue of the relation (Q5) used in [lr] and [BaHLR] is the Cuntz relation

$$(E3) \quad 1 = \sum_{m \in \Sigma} (u_m v) (u_m v)^*,$$

which is satisfied in the Cuntz-Pimsner algebra $\mathcal{O}(M_L)$ (see Proposition 4 below).

Proof. (Proof of Proposition 2) The Toeplitz algebra $\mathcal{T}(M_L)$ is generated by a universal representation $(i_{M_L}, i_{C(\mathbb{T}^d)})$. It is shown in [br, Corollary 3.3] that $\mathcal{T}(M_L)$ is generated by the range of $i_{C(\mathbb{T}^d)}$ and the single element $S := i_{M_L}(1)$, that $(i_{C(\mathbb{T}^d)}, S)$ is a Toeplitz-covariant representation in the sense of [br, Definition 3.1], and that $(\mathcal{T}(M_L), i_{C(\mathbb{T}^d)}, S)$ is universal for Toeplitz-covariant representations (ρ, V) satisfying

$$(TC1) \quad V\rho(a) = \rho(\alpha_A(a)) V, \text{ and}$$

$$(TC2) \quad V^* \rho(a) V = \rho(L(a)).$$

In our system $L(1) = 1$, and (TC2) implies that the operator V is an isometry.

The Stone-Weierstrass theorem implies that the characters $\gamma_m: z \mapsto z^m$ of \mathbb{T}^d span a dense $*$ -subalgebra of $C(\mathbb{T}^d)$, and a representation ρ of $C(\mathbb{T}^d)$ is completely determined by the unitary representation $u: m \mapsto \rho(\gamma_m)$ of \mathbb{Z}^d . One checks that $\alpha_A(\gamma_m) = \gamma_{Bm}$, so (TC1) is equivalent to (E1). We will complete the proof by showing that (TC2) is equivalent to (E2).

To see what (TC2) says about the u_m , we need to compute $L(\gamma_m)$. For any $f \in C(\mathbb{T}^d)$ and $z = e^{2\pi i x} \in \mathbb{T}^d$, we can compute $L(f)(z)$ by choosing one solution w_0 of $\sigma_A(w) = z$, such as $w_0 = e^{2\pi i A^{-1}x}$, and computing

$$L(f)(e^{2\pi i x}) = \frac{1}{|\det A|} \sum_{w \in \ker \sigma_A} f(w w_0),$$

and hence

$$L(\gamma_m)(e^{2\pi i x}) = \frac{1}{|\det A|} \gamma_m(e^{2\pi i A^{-1}x}) \sum_{w \in \ker \sigma_A} \gamma_m(w).$$

If $\gamma_m|_{\ker \sigma_A}$ is not the identity character, then $\{\gamma_m(w): w \in \ker \sigma_A\}$ is a non-trivial subgroup of \mathbb{T} , the sum is zero, and $L(\gamma_m) = 0$. So $L(\gamma_m) \neq 0 \iff \gamma_m \in (\ker \sigma_A)^\perp$, and for such m , $L(\gamma_m)(e^{2\pi i x}) = \gamma_m(e^{2\pi i A^{-1}x}) = \gamma_{B^{-1}m}(e^{2\pi i x})$. Now

$$\begin{aligned} \gamma_m \in (\ker \sigma_A)^\perp &\iff e^{2\pi i m^t A^{-1}n} = 1 \text{ for all } n \in \mathbb{Z}^d \\ &\iff e^{2\pi i (B^{-1}m)^t n} = 1 \text{ for all } n \in \mathbb{Z}^d \\ &\iff m \in B\mathbb{Z}^d. \end{aligned}$$

Thus

$$L(\gamma_m) = \begin{cases} 0 & \text{unless } m \in B\mathbb{Z}^d \\ \gamma_{B^{-1}m} & \text{if } m \in B\mathbb{Z}^d, \end{cases}$$

and using this we can see that (TC2) is equivalent to (E2).

For the last comment, recall from [br, §2] that the representation (ψ, π) corresponding to the Toeplitz-covariant representation (ρ, V) in the above argument is characterised by $\pi = \rho$ and $V = \psi(1)$, and that ρ satisfies $\rho(\gamma_m) = u_m$. \square

We now want an analogous presentation of $\mathcal{O}(M_L)$. To help keep things straight later, we write $\bar{u} := Q \circ u$ and $\bar{v} := Q(v)$.

Proposition 4. *Suppose that $A \in M_d(\mathbb{Z})$ has $\det A \neq 0$, and consider the Exel system $(C(\mathbb{T}^d), \alpha_A, L)$. Then the Cuntz-Pimsner algebra $\mathcal{O}(M_L)$ is the universal C^* -algebra generated by a unitary representation $\bar{u}: \mathbb{Z}^d \rightarrow U(\mathcal{O}(M_L))$ and an isometry $\bar{v} \in \mathcal{O}(M_L)$ satisfying (E1), (E2) and (E3).*

Proof. We need to prove that the unitary representation \bar{u} and the isometry \bar{v} satisfy (E1–3) and are universal for families satisfying these relations. They satisfy (E1) and (E2) because u and v do. To see that they satisfy (E3), note that the unitary u_m in Proposition 2 is $i_{C(\mathbb{T}^d)}(\gamma_m)$ and the isometry v is $i_{M_L}(1)$. We know from [ehr, Lemma 2.6] that $\{\gamma_m: m \in \Sigma\}$ is an orthonormal basis for M_L , so Lemma 2.5 of [ehr] says that a representation (ψ, π) is Cuntz-Pimsner covariant if and only if

$$1 = \sum_{m \in \Sigma} \psi(\gamma_m) \psi(\gamma_m)^*. \quad (2)$$

Since $\gamma_m = \gamma_m \cdot 1$ in M_L , we have

$$Q \circ i_{M_L}(\gamma_m) = Q \circ i_{C(\mathbb{T}^d)}(\gamma_m) Q \circ i_{M_L}(1) = Q(u_m) Q(v), \quad (3)$$

and Equation (2) for $(Q \circ i_{M_L}, Q \circ i_{C(\mathbb{T}^d)})$ reduces to (E3) for $Q \circ u$ and $Q(v)$.

Next suppose that U_m and V satisfy (E1), (E2) and (E3). Then Proposition 2 gives a representation (ψ, π) of M_L such that $U_m = \pi(\gamma_m)$ and $V = \psi(1)$, and, in view of (3), (E3) implies that ψ satisfies Equation (2). Thus (ψ, π) is Cuntz-Pimsner covariant, and hence factors through a representation of $\mathcal{O}(M_L)$, and this representation takes $\bar{u}_m = Q(u_m)$ to U_m and $\bar{v} = Q(v)$ to V because $\psi \times \pi$ takes u_m to U_m and v to V . \square

Next we want a convenient spanning family to do calculations with. Again, we are looking for something similar to what we used in [lr].

Lemma 5. *In $\mathcal{T}(M_L)$ we have*

$$\begin{aligned} (u_m \quad v^k \quad v^{*l} \quad u_n^*) \quad (u_p \quad v^i \quad v^{*j} \quad u_q^*) &= \\ \begin{cases} u_{m+B^{k-l}(p-n)} v^{k+i-l} v^{*j} u_q^* & \text{if } i \geq l \text{ and } p-n \in B^l \mathbb{Z}^d \\ u_m v^k v^{*(l+j-i)} u_{B^{j-i}(n-p)+q}^* & \text{if } i < l \text{ and } p-n \in B^i \mathbb{Z}^d \\ 0 & \text{otherwise.} \end{cases} \end{aligned} \quad (4)$$

Proof. We compute, using first (E2) and then (E1), to get

$$\begin{aligned} (u_m v^k v^{*l} u_n^*) (u_p v^i v^{*j} u_q^*) &= u_m v^k v^{*l} u_{p-n} v^i v^{*j} u_q^* \\ &= \begin{cases} u_m v^k u_{B^{-l}(p-n)} v^{i-l} v^{*j} u_q^* & \text{if } i \geq l \text{ and } p-n \in B^l \mathbb{Z}^d \\ u_m v^k v^{*(l-i)} u_{B^{-i}(n-p)}^* v^{*j} u_q^* & \text{if } i < l \text{ and } p-n \in B^i \mathbb{Z}^d \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} u_{m+B^{k-l}(p-n)} v^{k+i-l} v^{*j} u_q^* & \text{if } i \geq l \text{ and } p-n \in B^l \mathbb{Z}^d \\ u_m v^k v^{*(l+j-i)} u_{B^{j-i}(n-p)+q}^* & \text{if } i < l \text{ and } p-n \in B^i \mathbb{Z}^d \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

as required. \square

Corollary 6. *We have*

$$\mathcal{T}(M_L) = \overline{\text{span}}\{u_m v^k v^{*l} u_n^* : m, n \in \mathbb{Z}^d, k, l \in \mathbb{N}\}. \quad (5)$$

Proof. Equation (4) implies that $\text{span}\{u_m v^k v^{*l} u_n^*\}$ is a $*$ -algebra, and it contains all the generators of $\mathcal{T}(M_L)$. \square

Remark 7. *Since u is a unitary representation, we have $u_n^* = u_{-n}$, and the $*$ in u_n^* in (5) is technically redundant. We have retained the $*$ to emphasise the parallels between this situation and the one in [lr]. It also makes formulas more symmetric, and this sometimes simplifies calculations.*

4. A CHARACTERISATION OF KMS STATES

The gauge action $\gamma: \mathbb{T} \rightarrow \text{Aut}\mathcal{T}(M_L)$ is characterised by $\gamma_z(i_{M_L}(x)) = z i_{M_L}(x)$ and $\gamma_z(i_{C(\mathbb{T}^d)}(f)) = i_{C(\mathbb{T}^d)}(f)$, or equivalently by $\gamma_z(v) = z v$ and $\gamma_z(u_m) = u_m$. Our dynamics $\sigma: \mathbb{R} \rightarrow \text{Aut}\mathcal{T}(M_L)$ is defined in terms of the gauge action by $\sigma_t = \gamma_{e^{it}}$. Then we have

$$\sigma_t(u_m v^k v^{*l} u_n^*) = e^{it(k-l)} u_m v^k v^{*l} u_n^*,$$

which since $z \mapsto e^{iz(k-l)}$ is entire implies that the spanning elements $u_m v^k v^{*l} u_n^*$ are all analytic elements. Thus a state ϕ on $\mathcal{T}(M_L)$ is a KMS_β state for σ if and only if

$$\phi((u_m v^k v^{*l} u_n^*)(u_p v^i v^{*j} u_q^*)) = e^{-(k-l)\beta} \phi((u_p v^i v^{*j} u_q^*)(u_m v^k v^{*l} u_n^*)) \quad (6)$$

for all $m, n, p, q \in \mathbb{Z}^d$ and $i, j, k, l \in \mathbb{N}$.

The next result is an analogue of [lr, Lemma 8.3].

Proposition 8. *The system $(\mathcal{T}(M_L), \sigma)$ has no KMS_β states for $\beta < \log |\det A|$. For $\beta \geq \log |\det A|$, a state ϕ of $\mathcal{T}(M_L)$ is a KMS_β state if and only if*

$$\phi(u_m v^k v^{*l} u_n^*) = \begin{cases} 0 & \text{unless } k=l \text{ and } m-n \in B^k \mathbb{Z}^d \\ e^{-k\beta} \phi(u_{B^{-k}(m-n)}) & \text{if } k=l \text{ and } m-n \in B^k \mathbb{Z}^d. \end{cases} \quad (7)$$

Proof. Suppose that ϕ is a KMS_β state on $(\mathcal{T}(M_L), \sigma)$. Then for every $m \in \mathbb{Z}^d$, we have

$$\phi(u_m v v^* u_m^*) = \phi(v^* u_m^* \sigma_{i\beta}(u_m v)) = e^{-\beta} \psi(v^* u_m^* u_m v) = e^{-\beta} \psi(1) = e^{-\beta}.$$

Since $\{u_m v : m \in \Sigma\}$ is a Toeplitz-Cuntz family with $|\det A|$ elements, we have

$$1 = \phi(1) \geq \sum_{m \in \Sigma} \phi(u_m v v^* u_m^*) = |\det A| e^{-\beta}.$$

Thus $e^\beta \geq |\det A|$, and $\beta \geq \log |\det A|$.

Next we verify that ϕ satisfies (7). Applying the KMS condition twice gives

$$\phi(u_m v^k v^{*l} u_n^*) = e^{-k\beta} \phi(v^{*l} u_n^* u_m v^k) = e^{-(k-l)\beta} \phi(u_m v^k v^{*l} u_n^*).$$

Since $\beta \geq \log |\det A| > 0$, this implies that $\phi(u_m v^k v^{*l} u_n^*) = 0$ unless $k = l$. For $k = l$ we have

$$\phi(u_m v^k v^{*k} u_n^*) = e^{-k\beta} \phi(v^{*k} u_n^* u_m v^k) = e^{-k\beta} \phi(v^{*k} u_{m-n} v^k). \quad (8)$$

From k applications of (E2), we see that

$$v^{*k} u_{m-n} v^k = \begin{cases} 0 & \text{unless } m-n \in B^k \mathbb{Z}^d \\ u_{B^{-k}(m-n)} & \text{if } m-n \in B^k \mathbb{Z}^d, \end{cases}$$

and (8) implies (7).

Now we suppose that ϕ is a state of $\mathcal{T}(M_L)$ satisfying (7), and aim to prove that ϕ is a KMS_β state by verifying (6). There is a certain amount of symmetry to the two nonzero alternatives in formula (4), so we may as well assume that $i \geq l$ and $p-n \in B^l \mathbb{Z}^d$. Then (4) and (7) imply that $\phi((u_m u^k u^{*l} u_n^*)(u_p v^i v^{*j} u_q^*))$ is

$$\begin{cases} 0 & \text{unless } k+i-l=j \text{ and } m+B^{k-l}(p-n)-q \in B^j \mathbb{Z}^d \\ e^{-j\beta} \phi(u_{B^{-j}(m+B^{k-l}(p-n)-q)}) & \text{if } k+i-l=j \text{ and } m+B^{k-l}(p-n)-q \in B^j \mathbb{Z}^d. \end{cases}$$

The right-hand side of (6) also vanishes unless $k+i-l=j$, so we assume this from now on. Rewriting this equation as $k-j=l-i$ shows that our assumption $i \geq l$ is equivalent to $k \leq j$. Thus when we calculate the right-hand side of (6), the second alternative in (4) comes into play:

$$\begin{aligned} \phi((u_p \quad v^i \quad v^{*j} \quad u_q^*)(u_m \quad v^k \quad v^{*l} \quad u_n^*)) &= \\ \begin{cases} \phi(u_p v^i v^{*(j+l-k)} u_{B^{l-k}(q-m)+n}^*) & \text{if } q-m \in B^k \mathbb{Z}^d \\ 0 & \text{otherwise} \end{cases} \\ = \begin{cases} e^{-i\beta} \phi(u_{B^{-i}(p-B^{l-k}(q-m)-n)}) & \text{if } q-m \in B^k \mathbb{Z}^d \text{ and } p-B^{l-k}(q-m)-n \in B^i \mathbb{Z}^d \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Multiplying this by $e^{-(k-l)\beta}$ gives the right-hand side of (6), and in view of the equation $k+i-l=j$, it follows that the right-hand side of (6) is

$$\begin{cases} e^{-j\beta} \phi(u_{B^{-j}(p-B^{l-k}(q-m)-n)}) & \text{if } q-m \in B^k \mathbb{Z}^d \text{ and } p-B^{l-k}(q-m)-n \in B^j \mathbb{Z}^d \\ 0 & \text{otherwise.} \end{cases}$$

Since

$$\begin{aligned} B^{-i}(p-B^{l-k}(q-m)-n) &= B^{-i}(p-n) + B^{-(i+k-l)}(m-q) \\ &= B^{-j+k-l}(p-n) + B^{-j}(m-q), \end{aligned}$$

we have

$$e^{-j\beta} \phi(u_{B^{-j}(p-B^{l-k}(q-m)-n)}) = e^{-j\beta} \phi(u_{B^{-j}(m+B^{k-l}(p-n)-q)}),$$

so the numbers arising on the two sides of (6) are the same, and it remains to check that the conditions for nonvanishing are equivalent. So we need to check that

$$\begin{aligned} p - n \in B^l \mathbb{Z}^d \text{ and } m + B^{k-l}(p - n) - q \in B^j \mathbb{Z}^d \\ \iff q - m \in B^k \mathbb{Z}^d \text{ and } p - B^{l-k}(q - m) - n \in B^i \mathbb{Z}^d. \end{aligned}$$

Suppose that the first set of conditions holds. Then $q - m$ belongs to the coset $B^{(k-l)}(p - n) + B^j \mathbb{Z}^d$, which is contained in $B^k \mathbb{Z}^d$ because $k \leq j$ and $B^{-l}(p - n)$ is in \mathbb{Z}^d , and

$$p - B^{l-k}(q - m) - n = B^{l-k}(B^{k-l}(p - n) - (q - m))$$

belongs to $B^{l-k} B^j \mathbb{Z}^d = B^{l-k+j} \mathbb{Z}^d = B^i \mathbb{Z}^d$. So the forward implication holds, and similar arguments prove the converse. We have now proved (6), and thus ϕ is a KMS_β state. \square

5. KMS STATES FOR $\beta = \log |\det A|$.

We begin by showing that the Exel crossed product has very few KMS states.

Proposition 9. *If ϕ is a KMS_β state on $(\mathcal{O}(M_L), \sigma)$, then $\beta = \log |\det A|$.*

Proof. We compute using the relation (E3) and the KMS condition:

$$\begin{aligned} 1 = \phi(1) &= \phi\left(\sum_{m \in \Sigma} \bar{u}_m \bar{v} v^* \bar{u}_m^*\right) = \sum_{m \in \Sigma} \phi(\bar{u}_m \bar{v} v^* \bar{u}_m^*) \\ &= \sum_{m \in \Sigma} \phi(\bar{v}^* \bar{u}_m^* \sigma_{i\beta}(\bar{u}_m \bar{v})) = \sum_{m \in \Sigma} e^{-\beta} \phi(\bar{v}^* \bar{u}_m^* \bar{u}_m \bar{v}) \\ &= \sum_{m \in \Sigma} e^{-\beta} \phi(1) = N e^{-\beta}, \end{aligned}$$

since $|\Sigma| = |\mathbb{Z}^d / B\mathbb{Z}^d| = |\det B| = |\det A| = N$. \square

Lemma 10. *Every $\text{KMS}_{\log N}$ state of $(\mathcal{T}(M_L), \sigma)$ factors through the quotient map Q of $\mathcal{T}(M_L)$ onto $\mathcal{O}(M_L)$.*

Proof. Suppose that ϕ is a $\text{KMS}_{\log N}$ state of $(\mathcal{T}(M_L), \sigma)$. The formula (7) implies that $\phi(u_m v v^* u_m^*) = N^{-1}$, and hence

$$\phi\left(1 - \sum_{m \in \Sigma} u_m v v^* u_m^*\right) = 1 - N N^{-1} = 0.$$

Now the argument of [lr, Lemma 10.3] implies that ϕ vanishes on the ideal generated by $1 - \sum_{m \in \Sigma} u_m v v^* u_m^*$. But Proposition 4 says that this ideal is the kernel of Q , and the result follows. \square

The next result was first obtained by Ted Boey as an application of the general theory in [ln].

Theorem 11. *Suppose that $A \in M_d(\mathbb{Z})$ has $N := |\det A| \neq 0$. Then there is a $KMS_{\log N}$ state ϕ of $(\mathcal{O}(M_L), \sigma)$ such that*

$$\phi(\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*) = \begin{cases} 0 & \text{unless } k=l \text{ and } m=n \\ N^{-k} & \text{if } k=l \text{ and } m=n. \end{cases} \quad (9)$$

If A is a dilation matrix, then this is the only KMS state of $(\mathcal{O}(M_L), \sigma)$.

We will construct the state by factoring through an expectation onto the commutative subalgebra spanned by the range projections of the generators.

Lemma 12. *Suppose that $A \in M_d(\mathbb{Z})$ has nonzero determinant. Then there is an expectation E of $\mathcal{O}(M_L)$ onto*

$$\mathcal{O}(M_L)^\delta := \overline{\text{span}}\{\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^* : m \in \mathbb{Z}^d, k \in \mathbb{N}\} \quad (10)$$

such that

$$E(\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*) = \begin{cases} 0 & \text{unless } k=l \text{ and } m=n \\ \bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^* & \text{if } k=l \text{ and } m=n. \end{cases} \quad (11)$$

We prove this by averaging over a dual coaction, following a line of argument used in [lr1] and [lr] (and this explains the notation $\mathcal{O}(M_L)^\delta$). We will later give a second proof which avoids the use of coactions.

Proof. (First proof of Lemma 12) The Baumslag-Solitar group $\mathbb{Z}[B^{-1}] \rtimes \mathbb{Z}$ is the semidirect product of the additive subgroup $\mathbb{Z}[B^{-1}] := \bigcup_k B^{-k} \mathbb{Z}^d$ of \mathbb{Q} by the action of \mathbb{Z} by powers of B . We write ϵ for the canonical unitary representation of $\mathbb{Z}[B^{-1}] \rtimes \mathbb{Z}$ in $C^*(\mathbb{Z}[B^{-1}] \rtimes \mathbb{Z})$, so that in particular

$$\begin{aligned} \epsilon_{(0,1)} \epsilon_{(m,0)} &= \epsilon_{(Bm,1)} = \epsilon_{(Bm,0)} \epsilon_{(0,1)}, \text{ and} \\ \epsilon_{(0,1)}^* \epsilon_{(m,0)} \epsilon_{(0,1)} &= \epsilon_{(B^{-1}m,0)}. \end{aligned}$$

These identities imply that $U_m := \bar{u}_m \otimes \epsilon_{(m,0)}$ and $V := \bar{v} \otimes \epsilon_{(0,1)}$ satisfy the relations (E1), (E2) and (E3), and hence give a homomorphism $\delta := \pi_{U,V}$ of $\mathcal{O}(M_L)$ into $\mathcal{O}(M_L) \otimes C^*(\mathbb{Z}[B^{-1}] \rtimes \mathbb{Z})$. One can check on generators that δ is a coaction of $\mathbb{Z}[B^{-1}] \rtimes \mathbb{Z}$ on $\mathcal{O}(M_L)$. Since $\mathbb{Z}[B^{-1}] \rtimes \mathbb{Z}$ is amenable, averaging over this coaction gives an expectation E of $\mathcal{O}(M_L)$ onto the fixed-point algebra

$$\mathcal{O}(M_L)^\delta := \{a \in \mathcal{O}(M_L) : \delta(a) = a \otimes 1 = a \otimes \epsilon_{(0,0)}\}$$

(see [lr1, Lemma 6.5]). Since

$$\delta(\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*) = \bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^* \otimes \epsilon_{(m,k)(n,l)^{-1}},$$

$\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*$ belongs to the fixed-point algebra if and only if $(m,k) = (n,l)$, and thus E satisfies (11). Since E is norm-decreasing, and the $\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*$ span a dense subspace of $\mathcal{O}(M_L)$, (11) implies (10). \square

The coaction-free proof of Lemma 12 involves averaging twice over actions of abelian groups. Averaging over the gauge action $\gamma: \mathbb{T} \rightarrow \text{Aut } \mathcal{O}(M_L)$ gives an expectation E^γ onto the fixed-point algebra $\mathcal{O}(M_L)^\gamma$; since this expectation is continuous and kills elements $\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*$ with $k \neq l$, we have

$$\mathcal{O}(M_L)^\gamma = \overline{\text{span}}\{\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^*: m, n \in \mathbb{Z}^d, k \in \mathbb{N}\}. \quad (12)$$

For this proof of Lemma 12, we need to analyse the structure of $\mathcal{O}(M_L)^\gamma$, and since we'll use this analysis elsewhere in the proof of Theorem 11, we might as well do it properly now. As a point of notation, we write

$$\Sigma_k := \{\mu_1 + B\mu_2 + \cdots B^{k-1}\mu_k: \mu \in \Sigma^k\}, \quad (13)$$

and observe that Σ_k is a set of coset representatives for $\mathbb{Z}^d / B^k \mathbb{Z}^d$.

Proposition 13. (a) For each $k \geq 1$, we set

$$C_k := \overline{\text{span}}\{\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^*: m, n \in \mathbb{Z}^d\}.$$

Then the C_k are C^* -subalgebras of $\mathcal{O}(M_L)^\gamma$ satisfying $C_k \subset C_{k+1}$ and $\mathcal{O}(M_L)^\gamma = \bigcup_{k=1}^\infty C_k$.

(b) For each $k \geq 1$, $\{e_{m,n}^k := \bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^*: m, n \in \Sigma_k\}$ is a set of nonzero matrix units which spans a matrix algebra $M_{\Sigma_k}(\mathbb{C})$. The representation \bar{u} of \mathbb{Z}^d in $\mathcal{O}(M_L)$ maps $B^k \mathbb{Z}^d$ into C_k , and every $\bar{u}_{B^k m}$ belongs to the commutant of $M_{\Sigma_k}(\mathbb{C})$ in C_k .

(c) The inclusion ι_k of $M_{\Sigma_k}(\mathbb{C})$ in $\mathcal{O}(M_L)^\gamma$ and the integrated form $\pi_{\bar{u},k}$ of $\bar{u}|_{B^k \mathbb{Z}^d}$ give an isomorphism $\iota_k \otimes \pi_{\bar{u},k}$ of $M_{\Sigma_k}(\mathbb{C}) \otimes C^*(B^k \mathbb{Z}^d)$ onto C_k which carries $e_{m,n}^k \otimes \epsilon_{B^k p}$ into $\bar{u}_{m+B^k p} \bar{v}^k \bar{v}^{*k} \bar{u}_n^*$.

Proof. Calculations using the relations (E1) and (E2) show that

$$\begin{aligned} & (\bar{u}_m \quad \bar{v}^k \quad \bar{v}^{*k} \quad \bar{u}_n^*) \quad (\bar{u}_p \quad \bar{v}^k \quad \bar{v}^{*k} \quad \bar{u}_q^*) = \\ & \begin{cases} 0 & \text{unless } p - n \in B^k \mathbb{Z}^d \\ \bar{u}_{m+p-n} \bar{v}^k \bar{v}^{*k} \bar{u}_q^* & \text{if } p - n \in B^k \mathbb{Z}^d, \end{cases} \end{aligned} \quad (14)$$

which implies that C_k is a C^* -subalgebra. The Cuntz relation (E3) implies that

$$\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^* = \sum_{p \in \Sigma} \bar{u}_m \bar{v}^k (\bar{u}_p \bar{v} \bar{v}^* \bar{u}_p^*) \bar{v}^{*k} \bar{u}_n^* = \sum_{p \in \Sigma} \bar{u}_{m+B^k p} \bar{v}^{k+1} \bar{v}^{*(k+1)} \bar{u}_{n+B^k p}^*,$$

which in view of the definition of Σ_{k+1} implies that $C_k \subset C_{k+1}$. Now $\mathcal{O}(M_L)^\gamma = \bigcup_{k=1}^\infty C_k$ follows from (12), and we have proved (a).

Since $\{S_m := \bar{u}_m \bar{v}: m \in \Sigma\}$ is a Cuntz family, for each fixed k the products $\{S_\mu = S_{\mu_1} \cdots S_{\mu_k}: \mu \in \Sigma^k\}$ form a Cuntz family; since

$$S_\mu = (\bar{u}_{\mu_1} \bar{v}) (\bar{u}_{\mu_2} \bar{v}) \cdots (\bar{u}_{\mu_k} \bar{v}) = \bar{u}_{\mu_1 + B\mu_2 + \cdots B^{k-1}\mu_k} \bar{v}^k,$$

this Cuntz family is precisely $\{\bar{u}_m \bar{v}^k; m \in \Sigma_k\}$, and it follows that the $e_{m,n}^k$ are nonzero matrix units. The relation (E1) implies that $\bar{u}_{B^k p}$ commutes with every $e_{m,n}^k$, which gives (b).

Since the representation of $C^*(B^k \mathbb{Z}^d)$ in $C(\mathbb{T}^d)$ is faithful, and since $j_{C(\mathbb{T}^d)}: C(\mathbb{T}^d) \rightarrow C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N} = \mathcal{O}(M_L)$ is injective (by [br, Corollary 4.3], for example), $\pi_{\bar{u}, k}$ is injective, and hence so is the representation $\iota_k \otimes \pi_{\bar{u}, k}$ of $M_{\Sigma_k}(\mathbb{C}) \otimes C^*(B^k \mathbb{Z}^d) = M_{\Sigma_k}(C^*(B^k \mathbb{Z}^d))$. It is surjective because every $m \in \mathbb{Z}^d$ can be written uniquely as $m' + B^k m''$ for some $m' \in \Sigma_k$, and then

$$\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^* = \bar{u}_{B^k m''} (\bar{u}_{m'} \bar{v}^k \bar{v}^{*k} \bar{u}_{n'}^*) \bar{u}_{B^k n''}^*$$

is in C_k because each matrix unit and each $\bar{u}_{B^k m''}$ are. To see the last assertion about what $\iota_k \otimes \pi_{\bar{u}, k}$ does to $e_{m,n}^k \otimes \epsilon_{B^k p}$, recall that the representation $\phi \otimes \psi$ of a tensor product $C \otimes D$ coming from commuting representations ϕ of C and ψ of D takes $c \otimes d$ to $\phi(c) \psi(d) = \psi(d) \phi(c)$ (see [tfb, Theorem B.2], for example). \square

Corollary 14. *There is a continuous action τ of \mathbb{T}^d on $\mathcal{O}(M_L)^\gamma$ such that*

$$\tau_z(\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^*) = z^{m-n} \bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^* \text{ for } m, n \in \mathbb{Z}^d, k \in \mathbb{N}. \quad (15)$$

Proof. There is a continuous action η of \mathbb{T}^d on $M_{\Sigma_k}(\mathbb{C})$ such that $\eta_z(e_{m,n}^k) = z^{m-n} e_{m,n}^k$ — indeed, η_z is conjugation by the unitary $\sum_{m \in \Sigma_k} z^m e_{m,m}^k$. Lifting the dual action of $(B^k \mathbb{Z}^d)^\wedge = \mathbb{T}^d / (B^k \mathbb{Z}^d)^\perp$ to \mathbb{T}^d gives an action ζ of \mathbb{T}^d on $C^*(B^k \mathbb{Z}^d)$ which multiplies the generator $\epsilon_{B^k m}$ by $z^{B^k m}$. Pulling the action $\eta \otimes \zeta$ on $M_{\Sigma_k}(\mathbb{C}) \otimes C^*(B^k \mathbb{Z}^d)$ over to $\mathcal{O}(M_L)^\gamma$ under the isomorphism of Proposition 13(c) gives an action τ^k on C_k which satisfies (15) (for fixed k). A calculation using the Cuntz relation (E3) shows that the automorphisms τ_z^k combine to give an automorphism τ_z of $\bigcup_{k=1}^\infty C_k$, which is isometric because each τ_z^k is, and hence extends to an automorphism of $\mathcal{O}(M_L)^\gamma$. Continuity follows from the continuity of scalar multiplication. \square

Proof. (Second proof of Lemma 12) Averaging over the action τ of Corollary 14 gives an expectation E^τ of $\mathcal{O}(M_L)^\gamma$ onto $\mathcal{O}(M_L)^\delta$, and $E := E^\tau \circ E^\gamma$ has the required properties. \square

Proof. (Proof of existence in Theorem 11) The description of $\mathcal{O}(M_L)^\delta$ in (10) shows that each σ_t is the identity on $\mathcal{O}(M_L)^\delta$, so it suffices to find a trace τ on $\mathcal{O}(M_L)^\delta$ such that $\tau(\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^*) = N^{-k}$, and then Proposition 8 implies that $\tau \circ E$ is a $\text{KMS}_{\log N}$ state on $(\mathcal{O}(M_L), \sigma)$.

We can write each $m \in \mathbb{Z}^d$ uniquely as $m' + B^k m''$ for some $m' \in \Sigma_k$, and then part (b) of Proposition 13 implies that

$$\begin{aligned} \bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^* &= \bar{u}_{B^k m''} (\bar{u}_{m'} \bar{v}^k \bar{v}^{*k} \bar{u}_{m'}^*) \bar{u}_{B^k m''}^* \\ &= \bar{u}_{B^k m''} \bar{u}_{B^k m''}^* (\bar{u}_{m'} \bar{v}^k \bar{v}^{*k} \bar{u}_{m'}^*) \\ &= \bar{u}_{m'} \bar{v}^k \bar{v}^{*k} \bar{u}_{m'}^*. \end{aligned}$$

Now part (a) of Proposition 13 implies that

$$D_k := \text{span}\{\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^* : m \in \mathbb{Z}^d\} = \text{span}\{\bar{u}_{m'} \bar{v}^k \bar{v}^{*k} \bar{u}_{m'}^* : m' \in \Sigma_k\}$$

is a finite-dimensional commutative C^* -algebra, and that D_k has a normalised trace τ_k satisfying $\tau_k(\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^*) = N^{-k}$. The Cuntz relation (E3) implies that $D_k \subset D_{k+1}$, and the normalised traces τ_k combine to give a trace τ on $\mathcal{O}(M_L)^\delta = \bigcup_k D_k$ such that $\tau(\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^*) = N^{-k}$. Then, as foreshadowed above, $\phi := \tau \circ E$ is a $\text{KMS}_{\log N}$ state on $(\mathcal{O}(M_L), \sigma)$ satisfying (9). \square

For the proof of uniqueness, we need a standard fact about dilation matrices.

Lemma 15. *If B is an integer dilation matrix, then $\bigcap_{k=1}^\infty B^k \mathbb{Z}^d = \{0\}$.*

Proof. Suppose that $m \in \bigcap_{k=0}^\infty B^k \mathbb{Z}^d$. Then $B^{-k}m$ belongs to \mathbb{Z}^d for every k , and since we know from [ehr, Lemma 4.12], for example, that $\|B^{-k}m\| \rightarrow 0$ as $k \rightarrow \infty$, we must have $B^k m = 0$ for large k , and $m = 0$. \square

Proof. (Proof of uniqueness in Theorem 11) Suppose that ϕ is a KMS state of $(\mathcal{O}(M_L), \sigma)$. Proposition 9 implies that ϕ has inverse temperature $\beta = \log N$. We need to prove that ϕ satisfies (9), and comparing (9) with (7) (which we know holds with $e^{-k\beta} = N^{-k}$) shows that we need to prove that $\phi(\bar{u}_n) = 0$ for all nonzero n . So suppose $n \in \mathbb{Z}^d$ and $n \neq 0$. Lemma 15 implies that there is a smallest integer k such that n does not belong to $B^k \mathbb{Z}^d$. Then, recalling from the proof of existence that $\{\bar{u}_m \bar{v}^k : m \in \Sigma_k\}$ is a Cuntz family in $\mathcal{O}(M_L)$, we have

$$\phi(\bar{u}_n) = \phi\left(\bar{u}_n \sum_{m \in \Sigma_k} \bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_m^*\right) = \sum_{m \in \Sigma_k} \phi(\bar{u}_{n+m} \bar{v}^k \bar{v}^{*k} \bar{u}_m^*),$$

which vanishes by (7) because $(n+m) - m = n$ is not in $B^k \mathbb{Z}^d$ for every m . \square

It seems to be quite easy to find representations of $\mathcal{O}(M_L)$, and we describe an interesting one in the following example (which was one of our reasons for becoming interested in the C^* -algebras associated to dilation matrices in the first place [ehr]). However, it does not seem to be so easy to find natural Hilbert space representations of $\mathcal{O}(M_L)$ in which the $\text{KMS}_{\log|\det A|}$ state is a vector state.

Example 16. *The operators V and U_m on $L^2(\mathbb{T}^d)$ defined by*

$$(V\xi)(z) = \xi(\sigma_A(z)) \text{ and } (U_m \xi)(z) = z^m \xi(z)$$

satisfy (E1–3), and hence give a representation of $\mathcal{O}(M_L)$ on $L^2(\mathbb{T}^d)$. The Cuntz family $\{U_m V : m \in \Sigma\}$ is one of the sort studied by Bratteli and Jorgensen in the context of wavelets [bj], or more precisely, one of the more general sort studied in [ajln].

To make the connection, note that the characters $\{\gamma_n: n \in \Sigma\}$ form an orthonormal basis for the right Hilbert module M_L , or what is called in [aijln] a “filter bank for dilation by A ”. It is shown in [aijln, Proposition 2.2] that any filter bank $\{m_i: 0 \leq i < N\}$ gives rise to a Cuntz family $S_i := M(m_i) V$, where $M(f)$ is the operator of multiplication by $f \in C(\mathbb{T}^d)$. In the construction of wavelets, the more interesting filter banks are those in which m_0 is “low-pass”, which implies in particular that $m_0(1) = N^{1/2}$ and $m_i(1) = 0$ for $i > 0$ (see [aijln, Example 4.2]); the filters γ_n satisfy $|\gamma_n| \equiv 1$, and hence are “all-pass”.

Remark 17. Exel has previously studied KMS states on Exel crossed products [e2], and we now reconcile our result with his [e2, Proposition 9.2]. The situation in [e2] is more general than ours, but applies with $h = e 1$ and $E = \alpha \circ L$, which is easily seen to be an expectation of $C(\mathbb{T}^d)$ onto the range of α ; since our orthonormal basis for M_L is a quasi-basis, E has finite type with index $N := |\det A|$ (strictly speaking, $\text{ind} E$ is the element $N 1$ of $C(\mathbb{T}^d)$). Exel proved in [e2, Theorem 8.9] that there is an expectation $G: C(\mathbb{T}^d) \rtimes_{\alpha, L} \mathbb{N} \rightarrow j_{C(\mathbb{T}^d)}(C(\mathbb{T}^d))$ such that $G(\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*) = \delta_{k,l} N^{-k} \bar{u}_m \bar{u}_n^*$; in our situation, it is quite easy to check directly that G is given by first averaging over the gauge action γ , and then combining the expectations G_k on $C_k := \overline{\text{span}}\{\bar{u}_m \bar{v}^k \bar{v}^{*k} \bar{u}_n^*\}$ defined by $G_k(T) = N^{-k} \sum_{p \in \Sigma_k} \bar{u}_p T \bar{u}_p^*$ to get G on $(C(\mathbb{T}^d) \rtimes \mathbb{N})^\gamma = \bigcup_{k \geq 0} \overline{C_k}$ (see [ev, Corollary 7.5]). Then [e2, Proposition 9.2] implies that the KMS_β states on $C(\mathbb{T}^d) \rtimes_{\alpha, L} \mathbb{N}$ have the form $\phi \circ G$, where ϕ is a trace on $C(\mathbb{T}^d)$ satisfying $\phi(f) = e^{-\beta} N \phi(L(f))$ for $f \in C(\mathbb{T}^d)$.

Traces on $C(\mathbb{T}^d)$ are given by measures μ , and Exel’s condition says that μ satisfies

$$\int f \, d\mu = e^{-\beta} \int_{\mathbb{T}^d} \sum_{\sigma_A(w)=z} f(w) \, d\mu(z) \text{ for } f \in C(\mathbb{T}^d). \quad (16)$$

It follows from [aijln, Lemma 2.3], for example, that the Haar measure λ on \mathbb{T}^d satisfies (16) with $1 = e^{-\beta} N$, and since $\bar{u}_m \in C(\mathbb{T}^d) \rtimes \mathbb{N}$ is the image of the function z^n in $C(\mathbb{T}^d)$, the corresponding $\text{KMS}_{\log N}$ state ψ on $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ satisfies

$$\begin{aligned} \psi(\bar{u}_m \bar{v}^k \bar{v}^{*l} \bar{u}_n^*) &= \begin{cases} 0 & \text{unless } k=l \\ \int_{\mathbb{T}^d} N^{-k} z^m z^{-n} \, d\lambda(z) & \text{if } k=l \end{cases} \\ &= \begin{cases} 0 & \text{unless } k=l \text{ and } m=n \\ N^{-k} & \text{if } k=l \text{ and } m=n. \end{cases} \end{aligned}$$

Thus Exel’s result also gives the $\text{KMS}_{\log N}$ state described in Theorem 11, even though his state was obtained by factoring through a different expectation on $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$.

6. EXISTENCE OF KMS STATES FOR $\beta > \log |\det A|$.

Our goal here is to prove the existence of KMS_β states for $\beta > \log |\det A|$. Note that, when A is a dilation matrix, Lemma 15 implies that the sum on the right-hand side of (17) is finite.

Proposition 18. *Suppose that $A \in M_d(\mathbb{Z})$ satisfies $\det A \neq 0$ and that $\beta > \log |\det A|$. Then for each probability measure μ on \mathbb{T}^d , there is a KMS_β state $\psi = \psi_{\beta, \mu}$ of $(\mathcal{T}(M_L), \sigma)$ such that $\psi(u_m v^k v^{*l} u_n^*)$ vanishes unless $k = l$ and $m - n \in B^k \mathbb{Z}^d$, and*

$$\psi(u_m v^k v^{*l} u_n^*) = |\det A| e^{-\beta} \sum_{\{j \geq k : m-n \in B^j \mathbb{Z}^d\}} |\det A|^{j-k} e^{-j\beta} \int_{\mathbb{T}^d} z^{B^{-j}(m-n)} d\mu(z) \quad (17)$$

when $k = l$ and $m - n \in B^k \mathbb{Z}^d$.

We use the representation M of $C(\mathbb{T}^d)$ by multiplication operators on $L^2(\mathbb{T}^d, d\mu)$, and use the same notation for the corresponding unitary representation of \mathbb{Z}^d , so that $M_m := M(\gamma_m)$. For each $j \in \mathbb{N}$, we have a unitary representation $M \circ B^{-j}$ of the subgroup $B^j \mathbb{Z}^d$ of \mathbb{Z}^d , and we denote by \mathcal{H}_j the Hilbert space of the induced representation $\text{Ind}_{B^j \mathbb{Z}^d}^{\mathbb{Z}^d} M \circ B^{-j}$. Our state $\psi_{\beta, \mu}$ will be built from vector states for a representation π_μ of $\mathcal{T}(M_L)$ on $\mathcal{H}_\mu := \bigoplus_{j=0}^{\infty} \mathcal{H}_j$.

We will need to do some calculations in the Hilbert spaces \mathcal{H}_j , and for this it is convenient to use the sets Σ_j described in (13); for $g \in \mathbb{Z}^d / B^j \mathbb{Z}^d$, we write $c_j(g)$ for the element of Σ_j such that $c_j(g) \in g$. Then (from [tfb, page 296], for example) \mathcal{H}_j is the completion of the space

$\mathcal{V}_c := \{\xi: \mathbb{Z}^d \rightarrow L^2(\mathbb{T}^d, d\mu) \text{ such that } \xi(m-n) = M_{B^{-j}n}(\xi(m)) \text{ for } n \in B^j \mathbb{Z}^d\}$

in the inner product defined by

$$\begin{aligned} (\xi | \eta) &= \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} (\xi(c_j(g)) | \eta(c_j(g))) \\ &= \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} \int_{\mathbb{T}^d} \xi(c_j(g))(z) \overline{\eta(c_j(g))(z)} d\mu(z). \end{aligned}$$

(Although we have used the cross-section c_j to get a useful formula for the inner product, the translation condition on ξ and η means that this inner product does not depend on the choice of c_j .) Then the induced representation acts on \mathcal{H}_j by

$$((\text{Ind}_{B^j \mathbb{Z}^d}^{\mathbb{Z}^d} M \circ B^{-j})_m \xi)(n) = \xi(n - m).$$

We now take U to be the unitary representation of \mathbb{Z}^d on \mathcal{H}_μ defined by

$$U := \bigoplus_{j=0}^{\infty} (\text{Ind}_{B^j \mathbb{Z}^d}^{\mathbb{Z}^d} M \circ B^{-j}).$$

For each $j \geq 0$ and $\xi \in \mathcal{H}_j$, we define

$$(V_j \xi)(m) = \begin{cases} 0 & \text{unless } m \in B\mathbb{Z}^d \\ \xi(B^{-1}m) & \text{if } m \in B\mathbb{Z}^d; \end{cases}$$

a quick calculation shows that $V_j \xi$ belongs to \mathcal{H}_{j+1} . The V_j combine to give an isometry V on $\mathcal{H}_\mu = \bigoplus_j \mathcal{H}_j$, and the adjoint V^* is given on \mathcal{H}_{j+1} by the formula $(V^* \xi)(n) = \xi(Bn)$. Calculations show that the pair (U, V) satisfies (E1) and (E2), and hence there is a representation π_μ of $\mathcal{T}(M_L)$ on \mathcal{H}_μ such that $\pi_\mu(u_m) = U_m$ and $\pi_\mu(v) = V$.

We now let $e_{0,0}$ be the constant function 1 viewed as a unit vector in $\mathcal{H}_0 = L^2(\mathbb{T}^d, d\mu)$. For $j \in \mathbb{N}$ and $g \in \mathbb{Z}^d / B^j \mathbb{Z}^d$, we define $e_{j,g} := U_{c_j(g)} V^j e_{0,0}$, so that for each j ,

$$\{e_{j,g} : g \in \mathbb{Z}^d / B^j \mathbb{Z}^d\}$$

is an orthonormal set of $|\det B|^j = |\det A|^j$ vectors in \mathcal{H}_j . We view them as elements of \mathcal{H}_μ by adding 0s in the other summands. Inspired by the proof of [lr, Proposition 9.3], we define

$$\psi(T) := (1 - |\det A| e^{-\beta}) \sum_{j=0}^{\infty} \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} e^{-j\beta} (\pi_\mu(T) e_{j,g} | e_{j,g}).$$

Summing the geometric series $\sum_j (|\det A| e^{-\beta})^j$ shows that this series converges in norm in $\mathcal{T}(M_L)^*$, and that the sum is a state ψ of $\mathcal{T}(M_L)$.

Next we fix $m, n \in \mathbb{Z}^d$ and $k, l \in \mathbb{N}$, and verify the formula for $\psi(u_m v^k v^{*l} u_n^*)$. Then

$$V^{*l} U_n^* e_{j,g} = V^{*l} U_n^* U_{c_j(g)} V^j e_{0,0} = \begin{cases} 0 & \text{unless } l \leq j \\ V^{*l} U_{c_j(g)-n} V^j e_{0,0} & \text{if } l \leq j \end{cases}$$

belongs to \mathcal{H}_{j-l} , and hence

$$\begin{aligned} (\pi_\mu(u_m v^k v^{*l} u_n^*) e_{j,g} | e_{j,g}) &= (V^{*l} U_n^* e_{j,g} | V^{*k} U_m^* e_{j,g}) \\ &= \begin{cases} 0 & \text{unless } k = l \leq j \\ (V^{*k} U_{c_j(g)-n} V^j e_{0,0} | V^{*k} U_{c_j(g)-m} V^j e_{0,0}) & \text{if } k = l \leq j. \end{cases} \end{aligned}$$

We now recall that \mathcal{H}_{j-k} is the Hilbert space of the representation $\text{Ind}_{B^{j-k}\mathbb{Z}^d}^{\mathbb{Z}^d}(M \circ B^{j-k})$, and hence

$$\begin{aligned} &(V^{*k} U_{c_j(g)-n} V^j e_{0,0} | V^{*k} U_{c_j(g)-m} V^j e_{0,0}) \\ &= \sum_{h \in \mathbb{Z}^d / B^{j-k} \mathbb{Z}^d} (V^{*k} U_{c_j(g)-n} V^j e_{0,0}(c_{j-k}(h)) | V^{*k} U_{c_j(g)-m} V^j e_{0,0}(c_{j-k}(h))) \\ &= \sum_{h \in \mathbb{Z}^d / B^{j-k} \mathbb{Z}^d} (U_{c_j(g)-n} V^j e_{0,0}(B^k c_{j-k}(h)) | U_{c_j(g)-m} V^j e_{0,0}(B^k c_{j-k}(h))) \\ &= \sum_{h \in \mathbb{Z}^d / B^{j-k} \mathbb{Z}^d} (V^j e_{0,0}(B^k c_{j-k}(h) - c_j(g) + n) | V^j e_{0,0}(B^k c_{j-k}(h) - c_j(g) + m)). \end{aligned}$$

The h -summand vanishes unless both

$$B^k c_{j-k}(h) - c_j(g) + n \in B^j \mathbb{Z}^d \text{ and } B^k c_{j-k}(h) - c_j(g) + m \in B^j \mathbb{Z}^d. \quad (18)$$

As a function in the Hilbert space

$$\mathcal{H}_0 = \mathcal{H}(\text{Ind}_{\mathbb{Z}^d}^{\mathbb{Z}^d} M) = \{ \xi : \mathbb{Z}^d \rightarrow L^2(\mathbb{T}^d, d\mu) \text{ such that } \xi(-n) = M_n \xi(0) \},$$

$e_{0,0}$ satisfies $e_{0,0}(q)(z) = z^{-q}$, and $(V^j e_{0,0})(B^j q)(z) = e_{0,0}(q)(z) = z^{-q}$. Thus, when both criteria in (18) are satisfied, we have

$$\begin{aligned} (\pi_\mu(u_m v^k v^{*k} u_n^*) e_{j,g} \mid e_{j,g}) &= (V^{*k} U_{c_j(g)-n} V^j e_{0,0} \mid V^{*k} U_{c_j(g)-m} V^j e_{0,0}) \\ &= \int_{\mathbb{T}^d} z^{-B^{-j}(B^k c_{j-k}(h) - c_j(g) + n)} \overline{z^{-B^{-j}(B^k c_{j-k}(h) - c_j(g) + m)}} d\mu(z) \\ &= \int_{\mathbb{T}^d} z^{B^{-j}(m-n)} d\mu(z). \end{aligned}$$

(Notice that when (18) holds, we have $m - n \in B^j \mathbb{Z}^d$, so the last integral makes sense.) For each pair m, n such that $m - n$ is in $B^j \mathbb{Z}^d$, and each h in $\mathbb{Z}^d / B^{j-k} \mathbb{Z}^d$, there is exactly one g such that (18) holds. Thus, using (17) to view

$$\sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} e^{-j\beta} (\pi_\mu(u_m v^k v^{*k} u_n^*) e_{j,g} \mid e_{j,g}) \quad (19)$$

as a sum over $g \in \mathbb{Z}^d / B^j \mathbb{Z}^d$ and $h \in \mathbb{Z}^d / B^{j-k} \mathbb{Z}^d$, we find that (19) has exactly

$$|\mathbb{Z}^d / B^{j-k} \mathbb{Z}^d| = |\det A|^{j-k}$$

nonzero terms, each of which is equal to (18). Thus $\psi(u_m v^k v^{*k} u_n^*)$ vanishes unless $k=l$ and $m-n \in B^k \mathbb{Z}^d$, and then equals

$$(1 - |\det A| e^{-\beta}) \sum_{\{j \geq k : m-n \in B^j \mathbb{Z}^d\}} |\det A|^{j-k} e^{-j\beta} \int_{\mathbb{T}^d} z^{B^{-j}(m-n)} d\mu(z), \quad (20)$$

as stated in the Proposition.

We still need to prove that ψ is a KMS_β state, and we will do this using Proposition 8. So we need to compute $e^{-k\beta} \psi(u_{B^{-k}(m-n)})$ under the assumption that $m-n \in B^k \mathbb{Z}^d$. We have already done most of the work: the calculation (18) shows that

$$\begin{aligned} e^{-k\beta} \psi(u_{B^{-k}(m-n)}) &= e^{-k\beta} (1 - |\det A| e^{-\beta}) \sum_{j'=0}^{\infty} \sum_{g \in \mathbb{Z}^d / B^{j'} \mathbb{Z}^d} e^{-j'\beta} (U_{B^{-k}(m-n)} e_{j',g} \mid e_{j',g}) \\ &= e^{-k\beta} (1 - |\det A| e^{-\beta}) \sum_{\{j' : B^{-k}(m-n) \in B^{j'} \mathbb{Z}^d\}} \sum_{g \in \mathbb{Z}^d / B^{j'} \mathbb{Z}^d} e^{-j'\beta} \int_{\mathbb{T}^d} z^{B^{-j'} B^{-k}(m-n)} d\mu(z) \\ &= e^{-k\beta} (1 - |\det A| e^{-\beta}) \sum_{\{j' : B^{-k}(m-n) \in B^{j'} \mathbb{Z}^d\}} |\det A|^{j'} e^{-j'\beta} \int_{\mathbb{T}^d} z^{B^{-j'} B^{-k}(m-n)} d\mu(z), \end{aligned}$$

which reduces to (20) on writing $j = j' + k$. Thus Proposition 8 implies that ψ is a KMS_β state, and this completes the proof of Proposition 18.

7. PARAMETRISATION OF KMS_β STATES

Proposition 19. *Suppose that $A \in M_d(\mathbb{Z})$ has nonzero determinant and $\beta > \log |\det A|$. Then the map $\mu \mapsto \psi_{\beta, \mu}$ of Proposition 18 is an affine homeomorphism of the simplex $P(\mathbb{T}^d)$ of probability measures onto the simplex of KMS_β states for $(\mathcal{T}(M_L), \sigma)$.*

As in [lr, §10], the crux of the argument is a reconstruction formula which allows us to recover a KMS_β state from its “conditioning” ϕ_P to a corner $PT(M_L)P$. In the present situation, though, the projection

$$P := 1 - \sum_{g \in \mathbb{Z}^d / B\mathbb{Z}^d} u_{c(g)} v v^* u_{c(g)}^* = \prod_{g \in \mathbb{Z}^d / B\mathbb{Z}^d} (1 - u_{c(g)} v v^* u_{c(g)}^*)$$

belongs to $\mathcal{T}(M_L)$, so we don’t need to resort to spatial arguments to make sense of the conditioning: we can just define

$$\phi_P(a) = \frac{1}{1 - |\det A| e^{-\beta}} \phi(PaP),$$

and then since $\phi(u_{c(g)} v v^* u_{c(g)}^*) = e^{-\beta}$, the normalising factor ensures that ϕ_P is a state of $\mathcal{T}(M_L)$. We can now state our reconstruction formula.

Proposition 20. *Suppose that $\beta > \log |\det A|$, and that ϕ is a KMS_β state on $\mathcal{T}(M_L)$. Then for every $a \in \mathcal{T}(M_L)$ we have*

$$\begin{aligned} \phi(a) &= \lim_{n \rightarrow \infty} \left(1 - \sum_{j=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} e^{-j\beta} \phi_P(v^{*j} u_{c_j(g)}^* a u_{c_j(g)} v^j) \right). \end{aligned} \quad (21)$$

Convergence of the limit in Proposition 20 will follow from the following simple lemma:

Lemma 21. *Suppose that ϕ is a state of a unital C^* -algebra A , and that $\{p_n\}$ is a sequence of projections in A such that $\phi(p_n) \rightarrow 1$. Then $\phi(p_n a p_n) \rightarrow \phi(a)$ for every $a \in A$.*

Proof. We know that $\phi(1 - p_n) = 1 - \phi(p_n) \rightarrow 0$, so the Cauchy-Schwarz inequality for ϕ implies that $\phi(a(1 - p_n)) \rightarrow 0$ for all $a \in A$. Another application of the Cauchy-Schwarz inequality shows that $\phi(p_n a (1 - p_n)) \rightarrow 0$ also, so

$$\phi(a) - \phi(p_n a p_n) = \phi(a(1 - p_n)) + \phi((1 - p_n) a p_n) \rightarrow 0. \quad \square$$

When we apply Lemma 21, the projections p_n will be sums of the projections in the next proposition.

Proposition 22. *For $j \in \mathbb{N}$ and $g \in \mathbb{Z}^d / B^j \mathbb{Z}^d$ we define*

$$P_{j,g} := u_{c_j(g)} v^j P v^{*j} u_{c_j(g)}^*.$$

Then the $P_{j,g}$ are mutually orthogonal projections in $\mathcal{T}(M_L)$.

The proposition follows from the next lemma.

Lemma 23. *For each pair (j, g) and (l, h) we have*

$$P v^{*j} u_{c_j(g)}^* u_{c_j(h)} v^l P = \begin{cases} P & \text{if } j = l \text{ and } g = h \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If $j \neq l$, say $j < l$, then for $m \in \mathbb{Z}^d$ we have

$$P v^{*j} u_m v^l P = \begin{cases} P u_{B^{-j}m} v^{l-j} P & \text{if } m \in B^j \mathbb{Z}^d \\ 0 & \text{otherwise.} \end{cases}$$

Now every $n \in \mathbb{Z}^d$ (including $n = B^{-j}m$) has the form $n = c(n) + Bk$, so

$$P u_n v^{l-j} P = P u_{c(n)+Bk} v^{l-j-1} P = P u_{c(n)} v u_k v^{l-j-1} P,$$

which vanishes because P contains the factor $(1 - u_{c(n)} v v^* u_{c(n)}^*)$. So $P v^{*j} u_m v^l P$ vanishes when $j \neq l$, and for $j = l$, $P v^{*j} u_{c_j(h)-c_j(g)} v^j P$ vanishes unless $c_j(h) - c_j(g)$ belongs to $B^j \mathbb{Z}^d$, which occurs precisely when $g = h$ in $\mathbb{Z}^d / B^j \mathbb{Z}^d$. \square

Proof. (Proof of Proposition 20) We aim to apply Lemma 21 with

$$p_n := \sum_{j=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} P_{j,g},$$

which is a projection by Proposition 22. So we need to compute $\phi(p_n)$, which we do using the KMS condition:

$$\begin{aligned} \phi(p_n) &= \sum_{j=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} \phi(P_{j,g}) = \sum_{j=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} \phi(u_{c_j(g)} v^j P v^{*j} u_{c_j(g)}^*) \\ &= \sum_{j=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} e^{-j\beta} \phi(P v^{*j} u_{c_j(g)}^* u_{c_j(g)} v^j P) \\ &= \phi(P) \sum_{j=0}^n |\det A|^j e^{-j\beta} \quad (\text{by Lemma 23}) \\ &= (1 - |\det A| e^{-\beta}) \sum_{j=0}^n |\det A|^j e^{-j\beta}, \end{aligned}$$

which on summing the geometric series converges to 1 as $n \rightarrow \infty$. So Lemma 21 implies that for every $a \in \mathcal{T}(M_L)$, we have

$$\phi(a) = \lim_{n \rightarrow \infty} \sum_{j,l=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} \sum_{h \in \mathbb{Z}^d / B^l \mathbb{Z}^d} \phi(P_{j,g} a P_{l,h}).$$

Applying the KMS condition shows that this sum is

$$\lim_{n \rightarrow \infty} \sum_{j,l=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} \sum_{h \in \mathbb{Z}^d / B^l \mathbb{Z}^d} e^{-j\beta} \phi(P v^{*j} u_{c_j(g)}^* a u_{c_l(h)} v^l P v^{*l} u_{c_l(h)}^* (u_{c_j(g)} v^j P)),$$

and it follows from Lemma 23 that the summands are zero unless $j = l$ and $g = h$, in which case the right-hand factor $P v^{*l} u_{c_l(h)}^* u_{c_j(g)} v^j P$ collapses to P , and we recover the desired formula (21). \square

Proof. (Proof of Proposition 19) The formula (17) for $\psi_{\beta,\mu}$ shows that $\mu \mapsto \psi_{\beta,\mu}$ is affine and weak* continuous, and both sets of states are weak* compact, so it suffices to show that $\mu \mapsto \psi_{\beta,\mu}$ is surjective and one-to-one.

To see that $\mu \mapsto \psi_{\beta,\mu}$ is surjective, suppose that ϕ is a KMS_β state of $\mathcal{T}(M_L)$. On $C^*(u) = C(\mathbb{T}^d)$, the conditioned state ϕ_P is given by a probability measure μ ; we choose μ such that

$$\phi_P(u_m) = \int_{\mathbb{T}^d} z^m d\mu(z) \text{ for } m \in \mathbb{Z}^d,$$

and aim to prove that $\phi = \psi_{\beta,\mu}$. Since both states are KMS_β states, formula (7) shows that it suffices to check that $\phi(u_m) = \psi_{\beta,\mu}(u_m)$. Since \mathbb{Z}^d is abelian, the reconstruction formula (21) implies that

$$\begin{aligned} \phi(u_m) &= \lim_{n \rightarrow \infty} (1 - |\det A| e^{-\beta}) \sum_{j=0}^n \sum_{g \in \mathbb{Z}^d / B^j \mathbb{Z}^d} e^{-j\beta} \phi_P(v^{*j} u_m v^j) \\ &= \lim_{n \rightarrow \infty} (1 - |\det A| e^{-\beta}) \sum_{j=0}^n |\det A|^j e^{-j\beta} \phi_P(v^{*j} u_m v^j) \\ &= \lim_{n \rightarrow \infty} (1 - |\det A| e^{-\beta}) \sum_{\{j \leq n : m \in B^j \mathbb{Z}^d\}} |\det A|^j e^{-j\beta} \phi_P(u_{B^{-j}m}) \\ &= (1 - |\det A| e^{-\beta}) \sum_{\{j : m \in B^j \mathbb{Z}^d\}} |\det A|^j e^{-j\beta} \int_{\mathbb{T}^d} z^{B^{-j}m} d\mu(z), \end{aligned}$$

which by (17) is precisely $\psi_{\beta,\mu}(u_m)$. We have now proved surjectivity.

To see that our map is one-to-one, suppose that μ and ν are probability measures on \mathbb{T}^d and $\psi_{\beta,\mu} = \psi_{\beta,\nu}$. Write $M_\mu(n)$ for the n th moment $\int_{\mathbb{T}^d} z^n d\mu(z)$ of μ , and fix $m \in \mathbb{Z}^d$. Two applications of (17) show that

$$\begin{aligned} \sum_{\{j : m \in B^j \mathbb{Z}^d\}} |\det A|^j e^{-j\beta} M_\mu(B^{-j}m) &= \\ \sum_{\{j : m \in B^j \mathbb{Z}^d\}} |\det A|^j e^{-j\beta} M_\nu(B^{-j}m). \end{aligned} \quad (22)$$

The left-hand side of (22) can be rewritten as

$$\begin{aligned}
M_\mu(m) &+ \sum_{\{j : j > 0, m \in B^j \mathbb{Z}^d\}} |\det A|^j e^{-j\beta} M_\mu(B^{-j} m) \\
&= M_\mu(m) + |\det A| e^{-\beta} \sum_{\{j : j > 0, m \in B^j \mathbb{Z}^d\}} |\det A|^{j-1} e^{-(j-1)\beta} M_\mu(B^{-(j-1)} B^{-1} m) \\
&= M_\mu(m) + |\det A| e^{-\beta} \sum_{\{j' : B^{-1} m \in B^{j'} \mathbb{Z}^d\}} |\det A|^{j'} e^{-j'\beta} M_\mu(B^{-j'} B^{-1} m),
\end{aligned}$$

which by (17) is

$$\begin{cases} M_\mu(m) & \text{if } m \text{ is not in } B\mathbb{Z}^d \\ M_\mu(m) + |\det A| e^{-\beta} \psi_{\beta, \mu}(u_{B^{-1}m}) & \text{if } m \in B\mathbb{Z}^d. \end{cases}$$

If m is not in $B\mathbb{Z}^d$, then (22) says precisely that $M_\mu(m) = M_\nu(m)$; if $m \in B\mathbb{Z}^d$, then, since $\psi_{\beta, \mu}(u_{B^{-1}m}) = \psi_{\beta, \nu}(u_{B^{-1}m})$, subtracting $|\det A| e^{-\beta} \psi_{\beta, \mu}(u_{B^{-1}m})$ from both sides of (22) shows that $M_\mu(m) = M_\nu(m)$. Thus μ and ν have the same moments, and are therefore equal. \square

7.1. Limits of KMS states.

Proposition 19 describes all the KMS_β states for $\beta > \beta_c := \log |\det A|$, and Theorem 11 says there is exactly one KMS_{β_c} state when A is a dilation matrix. So it is natural to ask what we can say about the KMS_{β_c} states when A is not a dilation matrix. General results from [bra-rob] suggest that we might be able to find other KMS_{β_c} states by taking limits of KMS_β states as $\beta \rightarrow \beta_c$ from above.

Proposition 24. *Let $\mu \in P(\mathbb{T}^d)$. Then there is a decreasing sequence $\beta_n \rightarrow \beta_c$ such that $\{\psi_{\beta_n, \mu}\}$ converges weak* to a state ψ_μ , and then ψ_μ is a KMS_{β_c} state of $(T(M_L), \sigma)$.*

Proof. Choose any decreasing sequence converging to β_c , and the weak* compactness of the state space implies that there is a subsequence $\{\beta_n\}$ such that $\{\psi_{\beta_n, \mu}\}$ converges in the weak* topology. Now [bra-rob, Proposition 5.3.23] implies that the limit ψ_μ is a KMS_{β_c} state, at least when $\beta_c > 0$. When $\beta_c = 0$, [bra-rob, Proposition 5.3.23] only asserts that ψ_μ is a trace (because that is what being a KMS_0 state means in [bra-rob]). However, KMS_β states for $\beta > 0$ are σ -invariant, and hence so is the limit. Thus ψ_μ is a KMS_0 state in the sense we are using. \square

We now assume that A is not a dilation matrix, so that $\bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d$ could be bigger than $\{0\}$. Suppose $\beta > \beta_c$ and write $r = e^{-(\beta - \beta_c)}$. As in the last proof, we write $M_\mu(m)$ for the m th moment $\int_{\mathbb{T}^d} z^m d\mu(z)$. Rearranging (17) shows that $\psi_{\beta, \mu}(u_m v^k v^{*l} u_n^*)$ vanishes unless $k=l$ and $m-n \in B^k \mathbb{Z}^d$, and then equals

$$\sum_{\{j \geq 0 : m-n \in B^{j+k} \mathbb{Z}^d\}} e^{-k\beta} (1-r) r^j M_\mu(B^{-(j+k)}(m-n)). \quad (23)$$

So we want to compute the limit of (23) as $\beta \rightarrow \beta_c$, in which case $r \rightarrow 1$. If $m-n$ does not belong to $\bigcap_{j=0}^{\infty} B^{j+k} \mathbb{Z}^d = \bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d$, then the sum in (23) is finite, and since $(1-r)r^j \rightarrow 0$ as $r \rightarrow 1$ for each fixed j , (23) converges to 0 as $r \rightarrow 1$. So it remains for us to compute the limit of (23) when $m-n \in \bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d$. Unfortunately, this seems to be a fairly delicate matter (see Remark 25 below), and the best we can do is illustrate the issues with some examples.

- (a) If μ is normalised Haar measure on \mathbb{T}^d , then $M_\mu(0)=1$ and $M_\mu(m)=0$ for all other m . The series in (23) is identically zero unless $m=n$, and then is geometric; summing it shows that

$$\psi_{\beta, \mu}(u_m v^k v^{*l} u_n^*) = \begin{cases} 0 & \text{unless } k=l \text{ and } m=n \\ e^{-k\beta} & \text{if } k=l \text{ and } m=n. \end{cases}$$

Letting $\beta \rightarrow \beta_c$ gives the state described in Theorem 11.

- (b) If μ has the property that $M_\mu(m)=1$ for every $m \in \bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d$, then the series in (23) is geometric whenever $m-n \in \bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d$. Summing and letting $\beta \rightarrow \beta_c$ shows that the limit ψ_μ satisfies

$$\psi_\mu(u_m v^k v^{*l} u_n^*) = \begin{cases} 0 & \text{unless } k=l \text{ and } m-n \in \bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d \\ e^{-k\beta_c} & \text{if } k=l \text{ and } m-n \in \bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d. \end{cases}$$

- (c) The previous item (b) applies in particular to the point mass δ_1 at the identity $1=(1,1,\dots,1)$ of \mathbb{T}^d . This shows that the KMS_{β_c} state in Theorem 11 is unique if and only if $\bigcap_{j=0}^{\infty} B^j \mathbb{Z}^d = \{0\}$.
- (d) Consider the matrix $A = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$, for which $\bigcap_{j=0}^{\infty} B^j \mathbb{Z}^2 = \{0\} \times \mathbb{Z}$. Then item (b) applies to any measure of the form $\nu \times \delta_1$. Thus when A is not a dilation matrix, we expect there to be many KMS_{β_c} states besides the one in Theorem 11.
- (e) We wonder whether every KMS_{β_c} state is a limit of KMS_β states. It is trivially the case in our examples when $|\det A| > 1$, and in these examples it also works for $\beta_c=0$.

- (f) When $\beta_c = 0$, we have to be careful to distinguish between traces (the KMS_0 states in [bra-rob]) and the invariant traces (the KMS_0 states in [ped]). Certainly any limit of KMS_β states will be invariant, so the answer to the previous question is trivially false with the definition in [bra-rob] if the algebra has traces which are not invariant. We give an example where this happens in Remark 38.

Remark 25. *The obvious way to try to compute the limit of (23) as $r \rightarrow 1 -$ is to evaluate it term-by-term. This amounts to pulling $\lim_{r \rightarrow 1 -}$ through the infinite sum, and therefore requires the dominated convergence theorem. Write $m_j := M_\mu(B^{-(j+k)}(m - n))$. To apply the dominated convergence theorem, we need a convergent series $\sum_j a_j$ such that $0 \leq (1 - r)r^j |m_j| \leq a_j$ (and we need to consider a sequence $\{r_n\}$). We know $|m_j| \leq 1$. Calculus shows that $\max\{(1 - t)t^j : t \in [0, 1]\}$ occurs at $j/(j + 1)$. So the best general estimate seems to be*

$$(1 - r)r^j |m_j| \leq \left(1 - \frac{j}{j+1}\right) \left(\frac{j}{j+1}\right)^j = \frac{j^j}{(j+1)^{j+1}}.$$

Taking a_j to be the right-hand side and $b_j := 1/(j + 1)$, we have

$$\frac{b_j}{a_j} = \left(\frac{j+1}{j}\right)^j = \left(1 + \frac{1}{j}\right)^j \rightarrow e \text{ as } j \rightarrow \infty,$$

and the limit form of the comparison test implies that $\sum a_j$ diverges.

So pulling the limit through the sum seems to be a nontrivial matter. Of course, it is really just as well we can't do this, since we know that $\sum_{j=0}^{\infty} (1 - r)r^j = 1 \rightarrow 1$ as $r \rightarrow 1 -$, whereas the term-by-term calculation would give 0.

8. KMS_∞ AND GROUND STATES

Proposition 26. *Suppose that $A \in M_d(\mathbb{Z})$ has nonzero determinant. Then for every probability measure μ on \mathbb{T}^d , there is a KMS_∞ state $\psi_{\infty, \mu}$ on $(\mathcal{T}(M_L), \sigma)$ such that*

$$\psi_{\infty, \mu}(u_m v^k v^{*l} u_n^*) = \begin{cases} \int_{\mathbb{T}^d} z^{m-n} d\mu(z) & \text{if } k = l = 0 \\ 0 & \text{otherwise.} \end{cases}$$

Every ground state of $(\mathcal{T}(M_L), \sigma)$ has the form $\psi_{\infty, \mu}$, and is in particular a KMS_∞ state.

The proof of [lr, Lemma 8.4] gives the following characterisation of ground states.

Lemma 27. *A state ϕ of $\mathcal{T}(M_L)$ is a ground state for σ if and only if*

$$\phi_{\infty, \mu}(u_m v^k v^{*l} u_n^*) = \begin{cases} \phi(u_{m-n}) & \text{if } k = l = 0 \\ 0 & \text{otherwise.} \end{cases}$$

Proof. (Proof of Proposition 26) Choose a sequence $\{\beta_i\}$ such that $\beta_i \rightarrow \infty$; by passing to a subsequence, we may suppose that $\psi_{\beta_i, \mu}$ converges in the weak* topology to a state $\psi_{\infty, \mu}$, which is by definition a KMS_∞ state. Next we verify the formula for $\psi_{\infty, \mu}$. As $\beta \rightarrow \infty$, each summand in the right-hand side of (17) with $j > 0$ goes to zero. Thus as $i \rightarrow \infty$, we have

$$\psi_{\beta_i, \mu}(u_m v^k v^{*l} u_n^*) \rightarrow \begin{cases} \int_{\mathbb{T}^d} z^{m-n} d\mu(z) & \text{if } k=l=0 \\ 0 & \text{otherwise,} \end{cases}$$

and hence $\phi_{\infty, \mu}$ has the required form. (If A is not a dilation matrix, so that $\bigcap_{k=1}^\infty B^k \mathbb{Z}^d$ could contain nonzero elements, then the sum on the right-hand side of (17) could have infinitely many nonzero terms, and calculating the limit as $i \rightarrow \infty$ would require some analysis of the sort discussed in Remark 25.)

If ϕ is a ground state, then the restriction of ϕ to the range of $i_{C(\mathbb{T}^d)}$ is given by a probability measure μ , and then Lemma 27 implies that $\phi = \psi_{\infty, \mu}$. \square

9. THE TOEPLITZ ALGEBRA OF THE BAUMSLAG-SOLITAR SEMIGROUP

We fix an integer N with $N > 1$, and consider the additive group $\mathbb{Z}[N^{-1}]$ of rational numbers of the form $m N^{-l}$ for $m, l \in \mathbb{Z}$. The *Baumslag-Solitar group* is the semidirect product $\mathbb{Z}[N^{-1}] \rtimes \mathbb{Z}$ with

$$(r, k)(s, l) = (r + N^k s, k + l).$$

The semigroup semidirect product $\mathbb{N} \rtimes_N \mathbb{N}$ is a subsemigroup of $\mathbb{Z}[N^{-1}] \rtimes \mathbb{Z}$, and the pair $(\mathbb{Z}[N^{-1}] \rtimes \mathbb{Z}, \mathbb{N} \rtimes_N \mathbb{N})$ is closely related to the pair $(\mathbb{Q} \rtimes \mathbb{Q}_+, \mathbb{N} \rtimes \mathbb{N}^\times)$ studied in [lr]. Indeed, the map $(r, k) \mapsto (r, N^k)$ of $\mathbb{Z}[N^{-1}] \rtimes \mathbb{N}$ into $\mathbb{Q} \rtimes \mathbb{Q}_+$ carries $\mathbb{N} \rtimes_N \mathbb{N}$ into $\mathbb{N} \rtimes \mathbb{N}^\times$. The pair $(\mathbb{Z}[N^{-1}] \rtimes \mathbb{Z}, \mathbb{N} \rtimes_N \mathbb{N})$ is also quasi-lattice ordered in the sense of Nica [n]. (One way to see this is via the embedding of $\mathbb{Z}[N^{-1}] \rtimes \mathbb{N}$ in $\mathbb{Q} \rtimes \mathbb{Q}_+$: we just need to check that if $(r, k) \in \mathbb{Z}[N^{-1}] \rtimes \mathbb{N}$ and (r, N^k) has an upper bound in $\mathbb{N} \rtimes N^\mathbb{N}$, then the least upper bound constructed in [lr, Proposition 2.2] lies in $\mathbb{N} \rtimes N^\mathbb{N}$.) So $(\mathbb{Z}[N^{-1}] \rtimes \mathbb{Z}, \mathbb{N} \rtimes_N \mathbb{N})$ also has a Toeplitz algebra $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ which is universal for Nica covariant representations of $\mathbb{N} \rtimes_N \mathbb{N}$ [n, lr1]. The Toeplitz algebra $\mathcal{T}(C(\mathbb{T}), \alpha_N, L, \mathbb{N})$ is a quotient of $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ analogous to the additive boundary quotient of $\mathcal{T}(\mathbb{N} \rtimes \mathbb{N}^\times)$ studied in [BaHLR]. We now discuss the KMS states on $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$, following the analysis of [BaHLR, §4].

The Toeplitz algebra $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ is generated by two isometries $s = T_{(1,0)}$ and $v = T_{(0,1)}$, and an argument like that of [lr, §4] shows that $(\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N}), s, v)$ is universal among C^* -algebras generated by a pair of isometries S and V satisfying

$$(T1) \quad VS = S^N V,$$

$$(T4) \quad S^* V = S^{N-1} V S^*, \text{ and}$$

$$(T5) \quad V^* S^k V = 0 \text{ for } 1 \leq k < N.$$

We define $\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes_N \mathbb{N})$ to be the quotient of $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ by the extra relation $s s^* = 1$, $\mathcal{T}_{\text{mult}}(\mathbb{N} \rtimes_N \mathbb{N})$ to be the quotient by the relation $1 = \sum_{k=0}^{N-1} s^k v v^* s^{*k}$, and $\mathcal{T}_{\text{add, mult}}(\mathbb{N} \rtimes_N \mathbb{N})$ to be the quotient in which both extra relations hold, and which is therefore the analogue of Cuntz's $\mathcal{Q}_{(\text{mathbb{b}})} N$. Thus we have the following commutative diagram of quotient maps:

$$\begin{array}{ccc} & & \\ \text{\texttt{\textbackslash xymatrix}} & & \\ & & \\ \text{\texttt{\&\textbackslash TT(\textbackslash N\rtimes_N \textbackslash N)\textbackslash ar[d1]_{\textbackslash q_{\textbackslash add}}\textbackslash ar[dr]^{\textbackslash q_{\textbackslash mult}}\&\textbackslash}} & & \\ \text{\texttt{\&\textbackslash TT_{\textbackslash add}(\textbackslash N\rtimes_N \textbackslash N)\textbackslash ar[dr]\&\&\textbackslash TT_{\textbackslash mult}(\textbackslash N\rtimes_N \textbackslash N)\textbackslash ar[d1]\&\&}} & & \\ & & \\ \text{\texttt{\&\textbackslash TT_{\textbackslash add, \textbackslash ; \textbackslash mult}(\textbackslash N\rtimes_N \textbackslash N).\&\&}} & & \\ & & \end{array} \quad (24)$$

In $\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes_N \mathbb{N})$ the generator s becomes unitary, and (T4) is redundant. The unitary s generates a unitary representation $u: \mathbb{Z} \rightarrow U(\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes_N \mathbb{N}))$, and the relations (T1) and (T5) (taken together) are equivalent to (E1) and (E2) (taken together). Thus Proposition 2 implies that $\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes_N \mathbb{N})$ is our Toeplitz algebra $\mathcal{T}(M_L)$. Proposition 4 implies that $\mathcal{T}_{\text{add, mult}}(\mathbb{N} \rtimes_N \mathbb{N})$ is the quotient $\mathcal{O}(M_L)$ of $\mathcal{T}(M_L)$. (When $N = 2$, $\mathcal{O}(M_L)$ has been studied by Larsen and Li under the name \mathcal{Q}_2 ; see [LL, §3].)

Since the presentation of $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ is not affected by multiplying v by $z \in \mathbb{T}$, we can deduce from the presentation that there is an action $\gamma: \mathbb{T} \rightarrow \text{Aut} \mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ such that $\gamma_z(s) = s$ and $\gamma_z(v) = zv$. Inflating this action to \mathbb{R} gives a dynamics $\sigma: \mathbb{R} \rightarrow \text{Aut} \mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ such that $\sigma_t(s) = s$ and $\sigma_t(v) = e^{it} v$. This action leaves the kernels of the quotient maps in the diagram (24) invariant, and hence induces actions (still denoted by σ) on all three quotients. On $\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes_N \mathbb{N})$ and $\mathcal{T}_{\text{add, mult}}(\mathbb{N} \rtimes_N \mathbb{N})$ we recover the actions on $\mathcal{T}(M_L)$ and $\mathcal{O}(M_L)$ that we have been studying, in the case where A is the 1×1 matrix (N) and σ_A is the covering map $z \mapsto z^N$ of \mathbb{T} . So our results tell us about the KMS states of $(\mathcal{T}_{\text{add}}(\mathbb{N} \rtimes_N \mathbb{N}), \sigma)$ and $(\mathcal{T}_{\text{add, mult}}(\mathbb{N} \rtimes_N \mathbb{N}), \sigma)$.

Just as in [lr, Lemma 10.4], every KMS_β state of $(\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N}), \sigma)$ vanishes on the ideal generated by $1 - s s^*$, and hence comes from a KMS_β state of $\mathcal{T}(M_L)$. So we know all the KMS states of $(\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N}), \sigma)$. For ground states, though, there is a difference. As in [lr, Lemma 8.4] (or Lemma 27 above), a ground state of $(\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N}), \sigma)$ is determined by its values on $C^*(s)$, and we claim that the map $\phi \mapsto \phi|_{C^*(s)}$ is an affine homeomorphism of the set of ground states onto the state space of $C^*(s) \cong \mathcal{T}(\mathbb{N})$. Indeed, we can deduce this from [lr, Theorem 7.1(4)], since Theorem 3.7 of [lr1] implies that $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$ embeds as the subalgebra $C^*(s, v_N)$ of $\mathcal{T}(\mathbb{N} \rtimes \mathbb{N}^\times)$, and the homeomorphism $\phi \mapsto \phi|_{C^*(s)}$ factors through $C^*(s, v_N)$.

We can sum up these results by saying that the system $(\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N}), \sigma)$ has a phase transition at inverse temperature $\beta = \log N$, and a further phase transition (in the sense of Connes and Marcolli) at $\beta = \infty$. We believe that this is the simplest known system which exhibits both these phenomena. As for the system in [lr], the circular symmetry at $\beta = \log N$ which disappears for $\beta > \log N$ is not apparently realised by an action of \mathbb{T} on $\mathcal{T}(\mathbb{N} \rtimes_N \mathbb{N})$. In [lr], though, this circular symmetry persists for $\beta \in [1, 2]$, as a result of the more complicated convergence issues for the series representations of the normalising factors.

Remark 28. *Since we can view $\mathbb{N} \rtimes_N \mathbb{N}$ as a subsemigroup of $\mathbb{N} \rtimes \mathbb{N}^\times$, it might be more natural to use the dynamics satisfying $\sigma_t(v) = N^{it}v$. If we do this, then the phase transition will occur at $\beta = 1$.*

10. INTEGER MATRICES WITH DETERMINANT ± 1

When $A \in M_d(\mathbb{Z})$ has $|\det A| = 1$, the inverse A^{-1} has integer entries (as the cofactor formula shows), the map σ_A is a homeomorphism, and α_A is an automorphism. The inverse α_A^{-1} is then a transfer operator for α_A , so we have an Exel system $(C(\mathbb{T}^d), \alpha_A, \alpha_A^{-1})$, and this system has a Toeplitz algebra and an Exel crossed product. One would guess that these C^* -algebras must be related to the ordinary crossed product, and they are, but we have not seen this explicitly pointed out before.

Proposition 29. *Suppose that α is an automorphism of a unital C^* -algebra C . Then*

(a) *Exel's Toeplitz algebra $\mathcal{T}(C, \alpha, \alpha^{-1})$ is the universal C^* -algebra generated by an isometry v and a unital representation i_C of C satisfying $v i_C(c) = i_C(\alpha(c))v$, and*

(b) *the Exel crossed product $C \rtimes_{\alpha, \alpha^{-1}} \mathbb{N}$ is the universal C^* -algebra generated by a unitary u and a unital representation j_C of C satisfying $j_C(\alpha(c)) = u j_C(c) u^*$.*

Proof. We know from [br, §3] that $\mathcal{T}(C, \alpha, \alpha^{-1})$ is universal for Toeplitz-covariant representations (ρ, V) satisfying two relations called (TC1) and (TC2) (see page ?). As we observed earlier, plugging the identity 1 of C into (TC2) shows that V is an isometry. For our system (C, α, α^{-1}) , (TC1) implies (TC2):

$$V^* \rho(c) V = V^* \rho(\alpha(\alpha^{-1}(c))) V = V^* V \rho(\alpha^{-1}(c)) = \rho(\alpha^{-1}(c)),$$

and (a) follows.

To establish (b), notice first that $\phi(c) \in \mathcal{L}(M_{\alpha^{-1}})$ is the rank-one operator $\Theta_{c,1}$. The Cuntz-Pimsner algebra is generated by a universal Cuntz-Pimsner covariant representation $(j_{M_{\alpha^{-1}}}, j_C)$, and then the isometry v in part (a) is $v = j_{M_{\alpha^{-1}}}(1)$. Cuntz-Pimsner covariance says that

$$j_C(c) = (j_{M_{\alpha^{-1}}}, j_C)^{(1)}(\phi(c)) = (j_{M_{\alpha^{-1}}}, j_C)^{(1)}(\Theta_{c,1}) = j_{M_{\alpha^{-1}}}(c) j_{M_{\alpha^{-1}}}(1)^*;$$

since $c = c \cdot 1$, we have $j_{M_{\alpha^{-1}}}(c) = j_C(c) j_{M_{\alpha^{-1}}}(1)$, and Cuntz-Pimsner covariance is equivalent to $j_C(c) = j_C(c) v v^*$. This is equivalent to $v v^* = 1$, so v is unitary, and now $v j_C(c) = j_C(\alpha(c)) v$ is equivalent to $j_C(\alpha(c)) = v j_C(c) v^*$. \square

These universal properties immediately imply that our algebras are familiar objects:

Corollary 30. *Suppose that α is an automorphism of a unital C^* -algebra C . Then the Exel crossed product $C \rtimes_{\alpha, \alpha^{-1}} \mathbb{N}$ is the usual crossed product $C \rtimes_{\alpha} \mathbb{Z}$, and the Toeplitz algebra $\mathcal{T}(C, \alpha, \alpha^{-1})$ is the crossed product $C \rtimes_{\alpha} \mathbb{N}$ introduced and studied by Murphy [m]. In both cases, the gauge action of \mathbb{T} is the dual action of $\mathbb{T} = \hat{\mathbb{Z}}$.*

Remark 31. *Although $M_{\alpha^{-1}}$ is not the bimodule E considered by Pimsner in [p, Example (3), page 193], the two are isomorphic; indeed, $a \mapsto \alpha(a)$ is a Hilbert-bimodule isomorphism of E onto $M_{\alpha^{-1}}$. So the identity $C \rtimes_{\alpha, \alpha^{-1}} \mathbb{N} = C \rtimes_{\alpha} \mathbb{Z}$ also follows from the assertion in [p, Example (3)].*

We now return to the case of an integer matrix A with $|\det A| = 1$, where Corollary 30 identifies the Toeplitz algebra $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, \alpha_A^{-1})$ as a Murphy crossed product, and the Exel crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A, \alpha_A^{-1}} \mathbb{N}$ with the ordinary crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$. We will be working primarily with the crossed product, so it is worth observing that the generator \bar{v} is now unitary, and hence we can simplify our presentation: we view $C(\mathbb{T}^d) \rtimes_{\alpha} \mathbb{Z}$ as being generated by a unitary representation \bar{u} of \mathbb{Z}^d and a unitary \bar{v} satisfying $\bar{v} \bar{u}_m \bar{v}^* = \bar{u}_{Bm}$, and then

$$C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z} = \overline{\text{span}}\{\bar{u}_m \bar{v}^k : m \in \mathbb{Z}^d, k \in \mathbb{Z}\}.$$

We can if we wish make the further identification of the crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$ with the group algebra $C^*(\mathbb{Z}^d \rtimes_B \mathbb{Z})$ of the semidirect product (using Proposition 3.11 of [tfr2], for example).

As before, lifting the dual actions of \mathbb{T} gives actions σ of \mathbb{R} on $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, \alpha_A^{-1})$ and $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$ such that σ_t fixes the copies of $C(\mathbb{T}^d)$ and multiplies the additive generators by e^{it} . Proposition 18 describes the KMS_{β} states of $(\mathcal{T}(C(\mathbb{T}^d), \alpha_A, \alpha_A^{-1}), \sigma)$ for $\beta > \log |\det A| = 0$. Since $|\det B| = 1$, B is invertible over the integers, $B^j \mathbb{Z}^d = \mathbb{Z}^d$ for all j , and the series in (17) is infinite for every pair m, n . Thus for each $\mu \in P(\mathbb{T}^d)$ there is a KMS_{β} state $\psi_{\beta, \mu}$ on $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, \alpha_A^{-1})$ such that

$$\begin{aligned} \psi_{\beta, \mu} \left(\sum_{m, n} \langle u_m, v^k v^{*l} u_n^* \rangle \right) &= \\ \begin{cases} 0 & \text{unless } k = l \\ \sum_{j=k}^{\infty} (1 - e^{-\beta}) e^{-j\beta} M_{\mu}(B^{-j}(m - n)) & \text{if } k = l. \end{cases} \end{aligned} \quad (25)$$

Indeed, the proof of Proposition 18 simplifies substantially in this case: with $U = \bigoplus_{j=0}^{\infty} M \circ B^{-j}$ acting on $\bigoplus_{j=0}^{\infty} L^2(\mathbb{T}^d, d\mu)$, V the unilateral shift on the same direct sum, and e_j the constant function 1 in the j th summand and 0 elsewhere, we have

$$\psi_{\beta, \mu}(T) = \sum_{j=0}^{\infty} (1 - e^{-\beta}) e^{-j\beta} (\pi_{U, V}(T) e_j | e_j).$$

Proposition 18 also shows that all the KMS_0 states (that is, the invariant traces) on $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, \alpha_A^{-1})$ factor through traces of $C^*(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$. Since the uniqueness assertion in Theorem 11 does not apply, we might expect to find more than one.

Proposition 32. *Suppose that $A \in M_d(\mathbb{Z})$ has $|\det A| = 1$. If $\mu \in P(\mathbb{T}^d)$ satisfies $\sigma_A^* \mu = \mu$, then there is a σ -invariant trace ψ_μ on $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$ such that*

$$\psi_\mu(\bar{u}_m \bar{v}^k) = \begin{cases} 0 & \text{unless } k=0 \\ M_m(\mu) & \text{if } k=0, \end{cases} \quad (26)$$

and every σ -invariant trace on $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$ has this form.

Since the action σ of \mathbb{R} is inflated from the dual action $\hat{\alpha}_A$ of \mathbb{T} , a state is invariant for σ if and only if it is invariant for $\hat{\alpha}_A$. So the following standard lemma is useful.

Lemma 33. *Suppose that $\gamma: \mathbb{T} \rightarrow \text{Aut } D$ is a strongly continuous action on a unital C^* -algebra D and $E^\gamma: d \mapsto \int_{\mathbb{T}} \gamma_z(d) dz$ is the expectation onto the fixed-point algebra D^γ . Then a state ϕ of D is γ -invariant if and only if there is a state τ of D^γ such that $\phi = \tau \circ E^\gamma$.*

Proof. Suppose $\phi = \tau \circ E^\gamma$. Then the invariance of Haar measure implies that $E^\gamma \circ \gamma_z = E^\gamma$, and hence $\phi \circ \gamma_z = \tau \circ E^\gamma \circ \gamma_z = \tau \circ E^\gamma = \phi$, so ϕ is invariant. Conversely, if ϕ is invariant, then

$$\phi(d) = \int_{\mathbb{T}} \phi(d) dz = \int_{\mathbb{T}} \phi(\gamma_z(d)) dz = \phi\left(\int_{\mathbb{T}} \gamma_z(d) dz\right) = \phi \circ E^\gamma(d),$$

so $\phi = \phi|_{D^\gamma} \circ E^\gamma$; since $1 \in D^\gamma$, $\phi|_{D^\gamma}$ is a state. \square

Proof. (Proof of Proposition 32) With $\gamma = \hat{\alpha}_A$, the expectation E^γ is given by

$$E^\gamma(\bar{u}_m \bar{v}^k) = \begin{cases} 0 & \text{unless } k=0 \\ \bar{u}_m & \text{if } k=0. \end{cases}$$

It follows easily that if θ_μ is the state on $C(\mathbb{T}^d) = \overline{\text{span}}\{\bar{u}_m: m \in \mathbb{Z}^d\}$ given by integration against $\mu \in P(\mathbb{T}^d)$, then $\psi_\mu := \theta_\mu \circ E^\gamma$ satisfies (26). Lemma 33 implies that every invariant state of $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$ has this form. So we need to show that ψ_μ is a trace if and only if μ is invariant under σ_A^* .

We compute

$$\psi_\mu((\bar{u}_m \bar{v}^k)(\bar{u}_n \bar{v}^l)) = \begin{cases} \psi_\mu(\bar{u}_{m+B^k n}) & \text{if } k+l=0 \\ 0 & \text{otherwise,} \end{cases}$$

and similarly the other way round. Thus ψ_μ is a trace if and only if

$$\psi_\mu(\bar{u}_{m+B^{-l}n}) = \psi_\mu(\bar{u}_{n+B^l m}) = \psi_\mu(\bar{u}_{B^l(B^{-l}n+m)}) \text{ for all } l \in \mathbb{Z}, m, n \in \mathbb{Z}^d;$$

or, equivalently, if and only if $\psi_\mu(\bar{u}_m) = \psi_\mu(\bar{u}_{Bm})$ for all $m \in \mathbb{Z}^d$. But these are just the moments of μ , and a calculation shows that $M_{Bm}(\mu) = M_m(\sigma_A^* \mu)$. So we deduce that ψ_μ is a trace if and only if μ is invariant, as required. \square

Remark 34. When μ is invariant under σ_A , the moments $M_\mu(B^{-j}(m-n))$ appearing in (25) are all equal to $M_\mu(m-n)$. Thus the series on the right-hand side of (25) is geometric, and summing it shows that

$$\psi_{\beta, \mu}(u_m v^k v^{*l} u_n^*) = \begin{cases} 0 & \text{unless } k=l \\ e^{-k\beta} M_\mu(m-n) & \text{if } k=l, \end{cases}$$

which converges to $\psi_\mu(Q(u_m v^k v^{*l} u_n^*)) = \psi_\mu(\bar{u}_{m+B^{k-l}n} \bar{v}^{k-l})$ as $\beta \rightarrow 0$. In view of Proposition 24, this gives an alternative proof that ψ_μ is an invariant trace.

Corollary 35. When $|\det A| = 1$, $C(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$ has many invariant traces.

Proof. The homeomorphism σ_A has many finite orbits — indeed, the periodic points are dense in \mathbb{T}^d . (If the coordinates of $r \in \mathbb{Q}^d$ have common denominator N , then the denominators of all the coordinates in all the $A^n r$ divide N too, and the pigeon-hole principle implies that $e^{2\pi i r}$ is periodic.) But if $z \in \mathbb{T}^d$ has $\sigma_A^n(z) = z$, then $n^{-1} \sum_{j=0}^{n-1} \delta_{\sigma_A^j(z)}$ is an invariant measure. \square

Example 36. Consider $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, for which we have $\sigma_A(w, z) = (w, wz)$. Let λ denote Haar measure on \mathbb{T} and let ν be a probability measure on \mathbb{T} . Then the product measure $\mu = \nu \times \lambda$ is invariant for σ_A :

$$\begin{aligned} \int_{\mathbb{T}^2} f \circ \sigma_A(w, z) d\mu(w, z) &= \int_{\mathbb{T}} \int_{\mathbb{T}} f(w, wz) d\lambda(z) d\nu(w) \\ &= \int_{\mathbb{T}} \int_{\mathbb{T}} f(w, z) d\lambda(z) d\nu(w) = \int_{\mathbb{T}^2} f(w, z) d\mu(w, z). \end{aligned}$$

The moments of μ are given by

$$M_\mu(m) = \int_{\mathbb{T}} \int_{\mathbb{T}} w^{m_1} z^{m_2} d\lambda(z) d\nu(w) = \begin{cases} 0 & \text{unless } m_2 = 0 \\ M_\nu(m_1) & \text{if } m_2 = 0. \end{cases}$$

Thus Proposition 32 gives σ -invariant traces $\{\phi_\nu : \nu \in P(\mathbb{T})\}$ on $C^*(\mathbb{T}^d) \rtimes_{\alpha_A} \mathbb{Z}$ such that

$$\phi_\nu(\bar{u}_m \bar{v}^k) = \begin{cases} 0 & \text{unless } k=0 \text{ and } m_2=0 \\ M_\nu(m_1) & \text{if } k=0 \text{ and } m_2=0. \end{cases}$$

Example 37. When $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and $\mu = \lambda \times \nu$, we claim that the state $\psi_{\beta, \mu}$ described in (25) converges as $\beta \rightarrow 0+$ to the state of Theorem 11, which for $\beta = \log |\det A| = 0$ vanishes on $u_m v^k$ unless $(m, k) = (0, 0)$ and satisfies $\psi(u_0) = 1$. To see this, note that $B = A^t = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, so $B^{-j} = \begin{pmatrix} 1 & -j \\ 0 & 1 \end{pmatrix}$, and

$$M_{\lambda \times \nu}(B^{-j}(m-n)) = \int \int w^{(m_1-n_1)-j(m_2-n_2)} d\lambda(w) z^{m_2-n_2} d\nu(z),$$

which vanishes unless $m_1 - n_1 = j(m_2 - n_2)$. For $m = n$, we have $M_\mu(B^{-j}(m-n)) = 1$ for all j , and summing the series shows that $\psi_{\beta, \mu}(u_n v^k v^{*k} u_n^*) = e^{-k\beta} \rightarrow 1$ as $\beta \rightarrow 0$ for all k . If $m \neq n$ and $m-n$ does not have the form (li, i) , then $\psi_{\beta, \mu}(u_m v^k v^{*k} u_n^*) = 0$ for all k . If $m-n = (li, i)$, then $\psi_{\beta, \mu}(u_m v^k v^{*k} u_n^*)$ vanishes for $k > l$, and

$$\psi_{\beta, \mu}(u_m v^k v^{*k} u_n^*) = (1 - e^{-\beta}) e^{-l\beta} M_\nu(i) \rightarrow 0 \text{ as } \beta \rightarrow 0$$

for $k \leq l$. So whenever $m \neq n$, we have $\psi_{\beta, \mu}(u_m v^k v^{*k} u_n^*) \rightarrow 0$ as $\beta \rightarrow 0$, and the limit is the state ϕ of Theorem 11, as claimed. (Well, strictly speaking the limit is $\phi \circ Q$.)

Remark 38. When $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and $B = A^t = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, the map

$$(m, k) = ((m_1, m_2), k) \mapsto \begin{pmatrix} 1 & k & m_1 \\ 0 & 1 & m_2 \\ 0 & 0 & 1 \end{pmatrix} \quad (27)$$

is an isomorphism of $\mathbb{Z}^2 \rtimes_B \mathbb{Z}$ onto the integer Heisenberg group $H(\mathbb{Z})$. (The crux is that $B^k = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$.) For $\theta \in [0, 1]$, we view the rotation algebra \mathcal{A}_θ as the universal C^* -algebra generated by unitaries U, V satisfying $VU = e^{2\pi i \theta} UV$, and then the unitary representation $(m, k) \mapsto e^{2\pi i m_1 \theta} U^{m_2} V^k$ induces a surjection q_θ of $C^*(H(\mathbb{Z}))$ onto \mathcal{A}_θ . (Indeed, the quotients \mathcal{A}_θ are the fibres of a C^* -bundle over \mathbb{T} which has $C^*(H(\mathbb{Z}))$ as its algebra of continuous sections — see [e], [ap, §1] or [prae, Example 1.4].)

Every rotation algebra \mathcal{A}_θ has a trace τ_θ which kills $U^{m_2} V^k$ unless $m_2 = 0 = k$, and the composition $\tau_\theta \circ q_\theta$ is the invariant trace described in Example 36 for ν the point mass at $e^{2\pi i \theta}$. When θ is irrational, τ_θ is the only trace on \mathcal{A}_θ (see [dav, Proposition VI.1.3], for example). When θ is rational, \mathcal{A}_θ is a homogeneous C^* -algebra with spectrum \mathbb{T}^2 (by, for example, [dr, §2]), and has other traces which give non-invariant traces of $C^*(H(\mathbb{Z}))$. For example, the matrices $T = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $S = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ are unitary and satisfy $TS = -ST$, hence give a homomorphism $\pi_{S,T}: \mathcal{A}_{1/2} \rightarrow M_2(\mathbb{C})$, and composing with the usual normalised trace 2^{-1}tr gives a trace τ on $\mathcal{A}_{1/2}$ such that $\tau(U^{m_2} V^k) = 2^{-1}\text{tr}(S^{m_2} T^k)$. Since $T^2 = 1$, we have $\tau(V^2) = \tau(1) = 1$, and since $\sigma_t(v^2) = e^{2it} v^2$, $\tau(V^2) = 1$ implies that $\tau \circ q_{1/2}$ cannot be σ -invariant.

To explain where the other invariant traces in Example 36 come from, we examine the structure of $C^*(H(\mathbb{Z}))$ from another point of view. Consider the normal subgroup N of matrices (27) with $k = m_2 = 0$, which is the centre of $H(\mathbb{Z})$, and which has quotient $H(\mathbb{Z})/N$ isomorphic to \mathbb{Z}^2 via $(m, k) \mapsto (k, m_2)$. Applying Theorem 4.1 of [prae2] to N gives a realisation of $C^*(H(\mathbb{Z})) = \mathbb{C} \times_{*,1} H(\mathbb{Z})$ as a Busby-Smith twisted crossed product $C^*(N) \rtimes_{\beta, \omega} \mathbb{Z}^2$; identifying $C^*(N)$ with $C(\mathbb{T})$ and ploughing through the formulas in [prae2] shows that β is the identity and the cocycle $\omega: \mathbb{Z}^2 \rightarrow U(C(\mathbb{T})) = C(\mathbb{T}, \mathbb{T})$ is given by

$$\omega((k, m_2), (l, n_2))(z) = z^{kn_2}.$$

Averaging over the dual action of \mathbb{T}^2 gives an expectation $E^{\hat{\beta}}$ whose range is the fixed-point algebra $C(\mathbb{T}) \subset C(\mathbb{T}) \rtimes_{*, \omega} \mathbb{Z}^2$, which we can pull over to an expectation E on $C^*(H(\mathbb{Z}))$ such that

$$E(\bar{u}_m \bar{v}^k) = \begin{cases} 0 & \text{unless } k = 0 \text{ and } m_2 = 0 \\ \bar{u}_{m_1, 0} & \text{if } k = 0 \text{ and } m_2 = 0. \end{cases}$$

A direct calculation shows that E has the tracial property $E(ab) = E(ba)$ (E is a centre-valued trace on $C^*(H(\mathbb{Z}))$), and the isomorphism of $C^*(H(\mathbb{Z}))$ onto $C(\mathbb{T}) \rtimes_{*, \omega} \mathbb{Z}^2$ carries σ into $t \mapsto \hat{\beta}_{(e^{it}, 1)}$. Thus any state of the form $\phi \circ E$ is an invariant trace of $C^*(H(\mathbb{Z}))$. These are the traces described in Example 36.

BIBLIOGRAPHY

- [ap] J. Anderson and W. Paschke, *The rotation algebra*, Houston J. Math. **15** (1989), 1–26.
- [aijln] L.W. Baggett, N.S. Larsen, J.A. Packer, I. Raeburn and A. Ramsay, *Direct limits, multiresolution analyses, and wavelets*, J. Funct. Anal. **258** (2010), 2714–2738.
- [bos-con] J.-B. Bost and A. Connes, *Hecke algebras, type III factors and phase transitions with spontaneous symmetry breaking in number theory*, Selecta Math. (New Series) **1** (1995), 411–457.
- [bj] O. Bratteli and P.E.T. Jorgensen, *Isometries, shifts, Cuntz algebras and multiresolution analyses of scale N*, Integral Equations Operator Theory **28** (1997), 382–443.
- [bra-rob] O. Bratteli and D.W. Robinson, *Operator Algebras and Quantum Statistical Mechanics II*, Second Edition, Springer-Verlag, Berlin, 1997.
- [BaHLR] N. Brownlowe, A. an Huef, M. Laca and I. Raeburn, *Boundary quotients of the Toeplitz algebra of the affine semigroup over the natural numbers*, Ergodic Theory Dynam. Systems, to appear; arXiv:1009.3678.
- [br] N. Brownlowe and I. Raeburn, *\hat{E} Exel’s crossed product and relative Cuntz-Pimsner algebras*, Math. Proc. Cambridge Philos. Soc. **141** (2006), 497–508.
- [cm] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields, and Motives*, Colloquium Publications, vol. 55, Amer. Math. Soc., Providence, 2008.
- [cl] J. Crisp and M. Laca, *Boundary quotients and ideals of Toeplitz C^* -algebras of Artin groups*, J. Funct. Anal. **242** (2007), 127–156.
- [c2] J. Cuntz, *C^* -algebras associated with the $ax + b$ -semigroup over \mathbb{N}* , in: *K-Theory and Noncommutative Geometry* (Valladolid, 2006), European Math. Soc., Zürich, 2008, pages 201–215.

- [dav] K.R. Davidson, *C*-Algebras by Example*, Fields Institute Monographs, vol. 6, Amer. Math. Soc., Providence, 1996.
- [dr] S. Disney and I. Raeburn, *Homogeneous C*-algebras whose spectra are tori*, J. Aust. Math. Soc. (Ser. A) **38** (1985), 9–39.
- [e] G.A. Elliott, *On the K-theory of the C*-algebra generated by a projective representation of a torsion-free discrete group*, in: Operator Algebras and Group Representations, Vol I (Neptun, 1980), Monographs and Studies in Mathematics, vol. 17, Pitman, Boston, 1984, pages 157–184.
- [eva] D.E. Evans, *On \mathcal{O}_n* , Publ. Res. Inst. Math. Sci. **16** (1980), 915–927.
- [e1] R. Exel, *A new look at the crossed-product of a C*-algebra by an endomorphism*, Ergodic Theory Dynam. Systems **23** (2003), 1–18.
- [e2] R. Exel, *Crossed-products by finite index endomorphisms and KMS states*, J. Funct. Anal. **199** (2003), 153–188.
- [ehr] R. Exel, A. an Huef and I. Raeburn, *Purely infinite simple C*-algebras associated to integer dilation matrices*, Indiana Univ. Math. J., to appear; arXiv:1003.2097.
- [ev] R. Exel and A. Vershik, *C*-algebras of irreversible dynamical systems*, Canad. J. Math. **58** (2006), 39–63.
- [fr] N.J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999), 155–181.
- [laca98] M. Laca, *Semigroups of *-endomorphisms, Dirichlet series, and phase transitions*, J. Funct. Anal. **152** (1998), 330–378.
- [ln] M. Laca and S. Neshveyev, *KMS states of quasi-free dynamics on Pimsner algebras*, J. Funct. Anal. **211** (2004), 457–482.
- [lr1] M. Laca and I. Raeburn, *Semigroup crossed products and the Toeplitz algebras of nonabelian groups*, J. Funct. Anal. **139** (1996), 415–440.
- [lr] M. Laca and I. Raeburn, *Phase transition on the Toeplitz algebra of the affine semigroup over the natural numbers*, Adv. Math. **225** (2010), 643–688.
- [l] N. S. Larsen, *Crossed products by abelian semigroups via transfer operators*, Ergodic Theory Dynam. Systems **30** (2010), 1147–1164.
- [LL] N.S. Larsen and X. Li, *The 2-adic ring C*-algebra of the integers and its representations*, arXiv:1011.5622.
- [LR2] N.S. Larsen and I. Raeburn, *Projective multi-resolution analyses arising from direct limits of Hilbert modules*, Math. Scand. **100** (2007), 317–360.
- [m] G.J. Murphy, *Ordered groups and crossed products of C*-algebras*, Pacific J. Math. **148** (1991), 319–349.
- [n] A. Nica, *C*-algebras generated by isometries and Wiener-Hopf operators*, J. Operator Theory **27** (1992), 17–52.
- [prae2] J.A. Packer and I. Raeburn, *Twisted crossed products of C*-algebras*, Math. Proc. Cambridge Philos. Soc. **106** (1989), 293–311.
- [prae] J.A. Packer and I. Raeburn, *On the structure of twisted group C*-algebras*, Trans. Amer. Math. Soc. **334** (1992), 685–718.
- [pr] J.A. Packer and M.A. Rieffel, *Wavelet filter functions, the matrix completion problem, and projective modules over $C(\mathbb{T}^n)$* , J. Fourier Anal. Appl. **9** (2003), 101–116.
- [ped] G.K. Pedersen, *C*-Algebras and their Automorphism Groups*, London Math. Soc. Monographs, vol. 14, Academic Press, London, 1979.
- [p] M.V. Pimsner, *A class of C*-algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbb{Z}* , in: Free Probability Theory (Waterloo, 1995), Fields Institute Commun., vol. 12, Amer. Math. Soc., Providence, 1997, 189–212.
- [tfb] I. Raeburn and D.P. Williams, *Morita Equivalence and Continuous-Trace C*-Algebras*, Math. Surveys and Monographs, vol. 60, Amer. Math. Soc., Providence, 1998.
- [tfb^2] D.P. Williams, *Crossed Products of C*-Algebras*, Math. Surveys and Monographs, vol. 134, Amer. Math. Soc., Providence, 2007.

Proposition 39. *There is an expectation E of $C^*(H(\mathbb{Z}))$ onto $C^*(u_{(1,0)})$ such that*

$$E(u_{(m_1, m_2)} v^k) = \begin{cases} 0 & \text{unless } m_2 = k = 0 \\ u_{(m_1, 0)} & \text{if } m_2 = k = 0, \end{cases} \quad (28)$$

and we have $E(ab) = E(ba)$ for all $a, b \in C^*(H(\mathbb{Z}))$.

Proof. We consider the normal subgroup N of $H(\mathbb{Z})$ of matrices (27) with $k = m_2 = 0$, identify the quotient $H(\mathbb{Z})/N$ with \mathbb{Z}^2 via the map which sends the matrix (27) to (k, m_2) , and define $c: H(\mathbb{Z})/N \rightarrow H(\mathbb{Z})$ by

$$c(k, m_2) = \begin{pmatrix} 1 & k & 0 \\ 0 & 1 & m_2 \\ 0 & 0 & 1 \end{pmatrix}.$$

Applying the decomposition theorem [prae2, Theorem 4.1] to the twisted crossed product $C^*(H(\mathbb{Z})) = \mathbb{C} \times_{\cdot, 1} H(\mathbb{Z})$ with this data gives a twisted action (β, ω) of $\mathbb{Z}^2 = H(\mathbb{Z})/N$ on $\mathbb{C} \times_{\cdot, 1} N = C^*(N)$ and an isomorphism ϕ of $C^*(H(\mathbb{Z}))$ onto $C^*(N) \times_{\beta, \omega} \mathbb{Z}^2$. Since the original twisted action $(\cdot, 1)$ is trivial, and since N is central, the formulas in [prae2] collapse to give $\beta =$, and

$$\omega((k, m_2), (l, n_2)) = i_N(c(k, m_2) c(l, n_2) c(k + l, m_2 + n_2)^{-1}) = i_N\left(\begin{pmatrix} 1 & 0 & k n_2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}\right).$$

when we identify $C^*(N)$ with $C^*(\mathbb{Z})$ and $C(\mathbb{T})$ in the obvious way, $\omega: \mathbb{Z}^2 \times \mathbb{Z}^2 \rightarrow UC(\mathbb{T})$ is given by

$$\omega((k, m_2), (l, n_2))(z) = z^{k n_2}.$$

The formulas for the isomorphism ϕ (unfortunately buried in the proof of [prae2, Theorem 4.1]) also collapse, and we have

$$\phi(u_m v^k) = i_{C(\mathbb{T})}(\iota^{m_1}) i_{\mathbb{Z}^2}(k, m_2) = i_{\mathbb{Z}^2}(k, m_2) i_{C(\mathbb{T})}(\iota^{m_1}).$$

The crossed product $C(\mathbb{T}) \times_{\omega} \mathbb{Z}^2$ carries a natural dual action $\hat{\beta}$ of $\mathbb{T}^2 = (\mathbb{Z}^2)^\wedge$, which is characterised by

$$\hat{\beta}_{(w, z)}(i_{\mathbb{Z}^2}(k, m_2) i_{C(\mathbb{T})}(\iota^{m_1})) = w^k z^{m_2} i_{\mathbb{Z}^2}(k, m_2) i_{C(\mathbb{T})}(\iota^{m_1}).$$

Averaging over $\hat{\beta}$ gives an expectation of $C(\mathbb{T}) \times_{\omega} \mathbb{Z}^2$ onto $i_{C(\mathbb{T})}(C(\mathbb{T})) = i_{C(\mathbb{T})}(C^*(\iota))$, and pulling this back to $C^*(H(\mathbb{Z}))$ gives an expectation E onto $\phi^{-1}(i_{C(\mathbb{T})}(C^*(\iota))) = C^*(u_{(1,0)})$ satisfying (28). A direct computation with $a = u_m v^k$ and $b = u_n v^l$ gives the last assertion. \square